



Updated growth models for bigeye tuna (*Thunnus obesus*) in the Atlantic Ocean

Lynn Waterhouse^{a,*}, Lisa Ailloud^b, Riley Austin^c, Walter J. Golet^c, Ashley Pacicco^{d,2}, Allen H. Andrews^e, Khady Diouf^f, Yacine Ndiour^f, Kyne Krusic-Golub^g, Guelson da Silva^h, John M. Hoenigⁱ

^a John G. Shedd Aquarium, 1200 South Lake Shore Drive, Chicago, IL 60605, USA

^b National Marine Fisheries Service, Southeast Fisheries Science Center, 75 Virginia Beach Drive, Miami, FL 33149, USA.

^c The University of Maine – Gulf of Maine Research Institute, School of Marine Sciences, 350 Commercial Street, Portland, ME 04101, USA

^d Cooperative Institute for Marine and Atmospheric Studies, Rosenstiel School for Marine and Atmospheric Science, University of Miami, FL 33149, USA

^e University of Hawaii at Manoa, Department of Oceanography, 1000 Pope Road, Honolulu, HI 96822, USA

^f Institute fondamentale Afrique noire Cheikh Anta Diop - Corniche Ouest - Université Cheikh Anta Diop, BP 206 Dakar-Fann, Dakar, Senegal

^g Fish Ageing Services, 28 Swanston St, Queenscliff, 3225, VIC, Australia

^h Departamento de Ciências Animais, Universidade Federal Rural do Semiárido, Mossoro, Brazil

ⁱ Virginia Institute of Marine Science, College of William & Mary, P.O. Box 1346, Gloucester Point, VA 23062, USA

ARTICLE INFO

Handled by: Francisco J. Abascal

Keywords:

Validated age
Life history
Otolith
Scombridae
Growth model
Tagging

ABSTRACT

The International Commission for the Conservation of Atlantic Tunas (ICCAT) concluded the Atlantic Ocean tropical Tuna Tagging Programme (AOTTP) in 2021. This project had the objectives of enhancing food security, stimulating economic growth, and improving management through research on tropical tuna resources in the Atlantic Ocean, including bigeye tuna (*Thunnus obesus*). Here, we combine tagging data and otolith data from the AOTTP program, Panama City Lab and the Pelagic Fisheries Lab at the University of Maine with historical tagging data and otolith data from ICCAT and other sources to fit integrated growth models with the goal of providing the most complete growth curve (in terms of data inclusion and validation of age-at-length) for bigeye tuna in the Atlantic Ocean. Both Richards and von Bertalanffy growth models were fitted. A variety of models were fitted to subsets of the data to investigate the consistency of growth information. In all cases for the integrated model, the Richards and von Bertalanffy models were very similar with the von Bertalanffy model being preferred for parsimony. The preferred model, based on fit to old fish, was the von Bertalanffy curve based on length-age pair data from multiple sources. The addition of tagging data to create an integrated model showed patterns of lack of fit to both the tagging and otolith data suggesting conflict between the tagging and otolith data. The preferred model (length-age pair data only) gave the estimates: asymptotic length L_{∞} (fork length) equals 161.21 cm (95% bootstrap CI 154.39, 166.84), growth parameter K equals 0.392 yr^{-1} (95% bootstrap CI 0.355, 0.441), and the time-axis intercept t_0 equals -0.239 yr (95% bootstrap CI -0.306 , -0.175). For the best fitting integrated model, the asymptotic length L_{∞} (fork length, in cm) was estimated to be 185.78 (SD 6.298), the growth parameter K was 0.252 yr^{-1} (SD 0.014), and the time-axis intercept t_0 was -0.524 yr (SE 0.025). The value for asymptotic length L_{∞} from the integrated model was larger than the lengths of all the old fish in the sample whereas the value for the curve based on otoliths passes through the cloud of points for old fish.

* Corresponding author.

E-mail addresses: lwate@umn.edu (L. Waterhouse), lisa.ailloud@noaa.gov (L. Ailloud), riley.austin@maine.edu (R. Austin), walter.golet@maine.edu (W.J. Golet), ashley.pacicco@noaa.gov (A. Pacicco), aha7@hawaii.edu (A.H. Andrews), khady1.diouf@ucad.edu.sn (K. Diouf), yacine1.ndiour@ucad.edu.sn (Y. Ndiour), kyne.krusicgolub@fishageingservices.com (K. Krusic-Golub), guelson@ufersa.edu.br (G. da Silva), hoenig@vims.edu (J.M. Hoenig).

¹ Present address: U.S. Geological Survey, Minnesota Cooperative Fish and Wildlife Research Unit, University of Minnesota, Department of Fisheries, Wildlife, and Conservation Biology, 1980 Folwell Avenue, St. Paul, MN 55108, USA.

² Past address: Riverside Technology, Inc., Contractor to the National Marine Fisheries Service, Southeast Fisheries Science Center, 3500 Delwood Beach Road, Panama City, Florida 32408.

<https://doi.org/10.1016/j.fishres.2022.106317>

Received 31 May 2021; Received in revised form 21 March 2022; Accepted 24 March 2022

Available online 17 May 2022

0165-7836/© 2022 Published by Elsevier B.V.

1. Introduction

Bigeye tuna, *Thunnus obesus*, are distributed in the warm waters of the Atlantic, Pacific, and Indian Oceans where they are commercially and recreationally fished. They can grow up to 230 cm in length and weigh up to 250 kg (Collette and Nauen, 1983; Cayré et al., 1993). As a commercially valuable species fished in international waters by international parties, there is multi-national interest in keeping the fishery at a sustainable level. Accurately estimating the relationship between length and age provides critical information for assessment models. This paper focuses on the growth of bigeye tuna in the Atlantic Ocean.

Prior work for bigeye tuna in the Atlantic Ocean resulted in a variety of growth estimates, with values for the asymptotic maximum length, L_{∞} , ranging from ~200 to ~500 cm (Fig. 1, Appendix 1). These models relied on a variety of data sources to estimate growth including tagging, otoliths, spines, and length frequency data, all of which lacked old individuals and long-term tag recapture data. The 2021 International Commission for the Conservation of Atlantic Tunas (ICCAT) assessment of bigeye tuna utilized estimates from the Hallier et al. (2005) formulation of the Richards growth model, reparameterized using Schnute (1981) as required by the Stock Synthesis (SS3) assessment platform. The resulting parameters were: $L_{\infty} = 178.6$ cm fork length (FL), the growth parameter $K = 0.42 \text{ yr}^{-1}$ and the Richards coefficient $p = -0.00034$ (Anon, 2021).

New data collected on Atlantic bigeye tuna now allows the growth curve to be revisited. ICCAT concluded the Atlantic Ocean Tropical Tuna Tagging Programme (AOTTP) in 2021, a five-year program with the goal of tagging at least 120,000 tropical tunas with a variety of tag types (Beare et al., 2019). During this time, nearly 25,000 bigeye tuna were tagged and released with just over 5000 recoveries. Around the same time, laboratories across the Atlantic worked in collaboration to develop and validate age reading protocols for the species and increase the collection of hard parts for ageing purposes.

A goal of this work is therefore to estimate growth of Atlantic bigeye

tuna using all available data for the stock. Of particular interest was to combine multiple data sources (i.e., tagging data and otolith data, including validated ages) in order to develop the most comprehensive and up-to-date growth model for the species. This included tagging data from three different sources (AOTTP, the ICCAT historical database, and a study by Hallier et al., 2005) and otolith data from four different sources (AOTTP, Hallier et al., 2005, and age readings from the Pelagic Fisheries Lab (PFL) and the Panama City Lab (PCL) whose protocols have been validated using bomb radiocarbon dating (Andrews et al., 2020)).

With so many sources of information, it was clear from the outset that issues related to data quality would have to be addressed. Ailloud et al. (2014) found that ICCAT tagging data for Atlantic bluefin tuna contain useful information about growth rates if, and only if, the data are subjected to extensive quality control procedures. Such procedures have not been applied to the ICCAT bigeye tuna tagging data but have been applied to the tagging data from AOTTP (see Anon, 2021). In the present study, measurement error was estimated from short-term recapture data for the ICCAT and AOTTP tagging data as well as those of Hallier et al. (2005). It was also noted that different age reading protocols were used: Hallier et al. (2005) age estimates were based on daily growth rings while most other samples were aged using annual growth rings. While daily growth rings can provide accurate and precise age estimates in young fish, the procedure has been shown to progressively underestimate age for bigeye older than one year (Williams et al., 2013; Ailloud et al., 2019). As such, the data analysis considered several subsets of the full dataset.

In order to estimate growth simultaneously from tag-recapture data and otolith age-length data, the tag-recapture data must be modeled in a way that is consistent with age-length data (Francis 1988a,1988b; Laslett et al., 2002). That is, for age-length data, there is variability in length about age, so instead of modeling tag-recapture data as a function of length (i.e., using length increment data and times at liberty (Fabens 1965)) we model the lengths at release and recapture while treating the unknown age at release as a random variable (Francis et al., 2016; Aires-da-Silva et al., 2015; Eveson et al., 2004; Laslett et al., 2002). This modeling approach allows for the growth information from both sources of data to be modelled as a function of age, allowing for a common set of growth parameters to be derived.

2. Methods

2.1. Tagging Data

2.1.1. AOTTP Tagging Data

This analysis is based on AOTTP database version 'aottp_cisef 20210228'. Details of the AOTTP tagging program can be found in Beare et al. (2019). In the AOTTP database there are 24,252 releases of bigeye tuna (identified as bigeye in the release species code) representing 24,078 unique fish. Of those tagged fish there were 5018 recoveries (note, some of these represent fish recovered more than once from fish that were released post recovery).

Of the 5018 recoveries identified as bigeye during their release, 340 were identified during recovery as yellowfin (YFT), blackfin (BLF), little tunny (LTA), or skipjack (SKJ) (Table 1).

For the purpose of this analysis, we assume that the fish are bigeye if they were identified as bigeye during biological sampling. We also include fish that were identified as bigeye during release and recovery or




Fig. 1. Comparison of estimated growth curves for bigeye tuna from the literature and the current study. The curve by Hallier et al. was used in the most recent stock assessment by ICCAT. The two curves from the current study are without the age 1 and age 2 fish in the Pelagic Fisheries Lab dataset.

Table 1

Breakdown of species identifications during recovery of the 5018 identified as bigeye during release (BET = bigeye, BLF = Bluefin, LTA = Little Tunny, SKJ = Skipjack, YFT = Yellowfin, UNK = unknown).

BET	BLF	LTA	SKJ	YFT	UNK	Total
2243	1	1	22	316	2435	5018

Table 2

Sample size from AOTTP database for bigeye tuna data after each step in the data processing to ensure only appropriate pairs of data were used in the assessment work.

Initial number of releases in database	24,252
Initial number of paired (release-recapture) records in the database	5018
Justification for removal	# paired records remaining
<i>I – Initial data processing</i>	
Recovery length unknown or unable to convert to FL	4678
Missing time at liberty	4356
Time at liberty is negative	4280
Missing release length	4277
<i>II – Further exclusion criteria</i>	
Removed all records with time at liberty ≤ 97 days	1626
Removed outliers in growth	1592




Fig. 2. Plot of weekly growth (cm/week) versus time at liberty (weeks) based on straight fork length (FL) measurements at time of tagging and recapture from 4256 bigeye tuna in the AOTTP database. Only records for fish at liberty for up to 25 weeks are shown (maximum time at liberty is 161 weeks). The dashed vertical line is at 98 days. Due to the amount of data the circles have been made slightly transparent; circles that appear black (rather than grey) indicate multiple data points at this value.




Fig. 3. Histogram of lengths (cm) for the 1592 bigeye tuna used in this analysis at release (top) and recovery (bottom) from the AOTTP database.

bigeye during release and unknown during recovery. There are 4678 bigeye once this filter is applied (Table 2).

The release length type (relentype) and recovery length type (rclentype) were (straight) fork length (FL), blank, unknown (UNK), curved fork length (CFL), lower jaw to 1st dorsal (LD1), standard length (SL), or total length (TL). We retained fish which had length CFL (and




Fig. 4. Histogram of month of release (top) and recovery (bottom) for the 1592 bigeye tuna from the AOTTP database used in this analysis.

Table 3

Sample size from ICCAT database for bigeye Tuna data after each step in the data processing to ensure only appropriate pairs of data were used in the assessment work.

Initial number of releases in database	35,462
Initial number of paired (release-recapture) records in the database	7996
Justification for removal	# paired records remaining
<i>I – Initial data processing</i>	
Part of AOTTP database	2881
Recovery date is missing	2826
Release date is missing	2826
Release or recovery length is unknown	2108
Release or recovery measurement unit (e.g., FL or LJF) is unknown	45
Time at liberty is negative	44
<i>II – Further exclusion criteria</i>	
Removed all records with time at liberty ≤ 97 days	19
Removed any record that could overlap with Hallier	18
Removed growth outlier	17

converted them to FL, see Appendix 2 for details) and FL. We use the terms straight fork length and fork length interchangeably and distinct from curved fork length. There were 4356 bigeye tuna pairs left after this filter. After eliminating those fish with negative time at liberty we had 4280 pairs. There were 4227 remaining fish after those with missing lengths were removed; some had times at liberty up to 150 weeks.

Over short times at liberty the observed growth increments largely represent measurement error rather than somatic growth (Ailloud et al., 2014). We examined the distribution of unreasonable growth increments (negative weekly growth) as a function of time at liberty to determine a threshold time at liberty at which measurement errors are minimal while retaining as great a sample size as possible. In order to match what was done across other datasets, 98 days was used as the cutoff for determining the time at liberty that represents real growth rather than measurement error. This left 1626 records (Table 2). In an attempt to eliminate outliers due to data entry and measurement errors, we removed records with the fastest and slowest 1% absolute growth per day (i.e., below the 0.01 and greater than the 0.99 quantiles). This resulted in 1592 records for analysis.

The resulting dataset for bigeye consisted of 1592 records with lengths at tagging ranging from 33 cm FL to 140 cm FL, lengths at recapture ranging from 28 cm FL to 150 cm FL (Fig. 3) and times at liberty ranging from 98 to 1127 days (median = 239 days). The releases peaked in March and then July to November, while the majority of the recoveries occurred May to August with a peak in July (Fig. 4).




Fig. 5. Histogram of month of release (top) and recovery (bottom) for the 18 bigeye tuna used in this analysis (prior to the removal of the growth outlier), from the ICCAT database.




Fig. 8. Frequency of recovery fork lengths (cm), top panel, and release lengths (cm), bottom panel, for bigeye tuna from the tagging data of Hallier et al.




Fig. 6. Histogram of lengths (cm) for the 18 bigeye tuna used in this analysis at release (top) and recovery (bottom) from the ICCAT database. The fish with recovery length of 203 cm is an apparent outlier and was discarded (see Fig. 10).




Fig. 7. Growth per week (recorded growth divided by weeks at liberty) versus time at liberty for bigeye tuna from the ICCAT database. The dashed vertical line denotes the 98 day time at liberty cutoff used in this study.

2.2. ICCAT tagging data

The ICCAT tagging database (ICCAT Secretariat (n.d.)) begins in 1960 for bigeye tuna and has releases or recaptures for 54 years (through 2020, which overlaps with the AOTTP tagging program). The




Fig. 9. Growth per week (recorded growth divided by weeks at liberty) versus time at liberty for bigeye tuna from the tagging data of Hallier et al. The dashed vertical line denotes the 98 day time at liberty cutoff used in this study.

tagging database has a total of 35,462 releases and 7996 recoveries, although many of these (24,212 releases and 5115 recoveries) are part of the AOTTP database (Table 3). A further 55 records have unknown release years and 115 records had missing recovery years. There are 2108 pairs of release and recoveries with known dates of release and recovery and known lengths at release and recovery.

Of the 2108 pairs of release/recovery, only 45 had both pairs with known measurement unit (either FL or LJF), and the rest were unknown. We treated LJF (lower jaw fork length) as equivalent to FL and thus all 45 lengths were retained. After removing those fish with negative time at liberty we had 44 pairs.

The majority of the tagged bigeye from the ICCAT database were tagged in the months of July and August with recoveries throughout the year, but with a peak in August (Fig. 5). The release FL ranged from 56 to 129 cm and the recovery FL ranged from 12.5 to 203 cm (Fig. 6).

In order to avoid including records where observed growth rates most likely reflect measurement error or tagging effects, only data for fish at liberty > 97 days were retained, leaving 19 data pairs to analyze (Fig. 7). Of these 19 pairs, 16 had lengths that were measured at both release and recovery; 2 pairs had unknown (either estimated or measured) measurement type at release and recovery; and 1 pair had estimated measurement type at release and unknown at recovery. One additional record was removed to avoid overlapping with the Hallier et al. (2005) data (discussed below), as those data occur in the ICCAT database, but we were unable to uniquely identify the records. The one removal was a release from 2000 and this overlaps with the time frame of the Hallier data (releases in 1994–2000). Another record has an




Fig. 10. Plot of all the length-age pair data (excluding fish greater than one year of age in the dataset of Hallier et al., 2005). Blue symbols represent data from Panama City Lab; cyan = ages from daily ring counts by Hallier et al. (2005), green symbols = AOTTP readings of annual rings in otoliths, magenta = data from Pelagic Fisheries Lab (PFL). Integer ages > 1 have been jittered to reduce overprinting. Four fitted von Bertalanffy curves are shown which differ in whether age 1 and age 2 fish from the PFL dataset are included and whether tagging data is used in addition to the otolith data.

unreasonable growth trajectory (see Fig. 10) and was removed. This resulted in 17 usable records from the ICCAT database.

2.3. Tagging data from Hallier et al. (2005)

We obtained data from a previous study in the eastern Atlantic Ocean (Hallier et al., 2005) that included 625 bigeye tagged and recaptured with FL ranging from 37 to 124 cm (Fig. 8). When 98 days is used as the time at liberty cutoff (to be consistent with AOTTP and ICCAT tagging data), 146 bigeye release and recoveries remain (Fig. 9). Originally Hallier used 14 days as the cutoff.

2.4. Otolith data

The otolith data comprised samples aged using two different reading protocols: one based on daily increment counts and the other on annual ring counts. Results from blind counts of micro-increments in chemically marked bigeye tuna have found that micro-increment counts tend to underestimate true times at liberty (Ailloud et al., 2019; Farley et al., 2020), indicating that daily counts are likely to underestimate true age. Additional work by Williams et al. (2013) has shown that age counts of presumed daily growth increments can lead to an underestimation of age in fish older than 1 year (Williams et al., 2013). As such, for all hard part data utilized in this analysis, daily readings were restricted to bigeye less than 1 year of age. The ageing protocol based on annual ring counts (Allman et al., 2020) that was used for all samples described below has been validated through bomb radiocarbon dating (Andrews et al., 2020) and preliminary results from AOTTP fish marked with

oxytetracycline support the hypothesis that the larger growth increments are deposited on an annual basis (Ailloud et al., 2019).

2.5. AOTTP otolith data

A total of 63 pairs of otolith age and length was obtained through the AOTTP from fish sampled across a large area of the Atlantic Ocean. Twenty-six of those samples were read by a single expert using annual growth increment counts, while the remaining 37 (the 'reference collection') were aged by three independent readers who counted daily rings (Fig. 10; Beare et al., 2019).

Whole otoliths were imaged and weighed prior to processing. Each core was marked prior to embedding in embedding in polyplex clear ortho casting resin (Allnex®). Transverse sections were cut through the center of each otolith using an Isomet 1000 low-speed saw with diamond edged wafering blades. Sections were then mounted on microscope slides (76.2 × 25.4 mm) using thermoplastic resin (Cystalbond 509 ©) with the side of the section furthest away from the core facing up. Each section was ground to a thickness between 320 and 350 µm using wet/dry sandpaper (800 and 1200 grit), lubricated with distilled water. A small drop of microscope immersion oil (Cargille © -TYPE A) was added prior to imaging to help clear up the ground surface of the otolith and aid in the imaging process.

Otolith sections were imaged under transmitted light. Annual ages were assigned to individuals based on the number of fully formed opaque zones (i.e., presence of translucent otolith material between the outer edge of the last opaque zone and the otolith margin). All age readings were made without knowledge of fish size, otolith weight, sex, location of capture or time at liberty. Methods for the annual age reading followed those developed for other tuna species (Farley et al., 2013, 2006; Gunn et al., 2008; Lang et al., 2017). Ageing protocols developed for Atlantic bluefin tuna (Neilson and Campana, 2008; Rodríguez-Marín et al., 2004, 2007; Secor et al., 2014) were also used as a basis to aid interpretation of what may constitute an annual growth zone in Atlantic tunas.

Micro-increment counts were conducted at various magnifications ranging between 400 and 1000x. The method for the interpretation of the microstructure was consistent with those methods published for reading transverse sections (Lehodey and Leroy, 1999; Sardenne et al., 2015; Shuford et al., 2007). After a count of between 150 and 180 the internal micro-structure becomes increasingly difficult to interpret. For subsections of the otolith where increments were either difficult to interpret or not present, an interpolation method based on the zone pattern immediately before and after the difficult area was applied.

2.5.1. Otolith data from pelagic fisheries lab

A total of 229 sets of otoliths was extracted from bigeye tuna landed by commercial pelagic longline vessels and recreational rod and reel fisheries along the east coast of the United States between June and November of 2018–2020. Catch locations include the Gulf Stream, along the continental shelf, and slope canyons from Cape Hatteras to the Hague Line. FL (cm) of individuals sampled from recreational fisheries and CFL from commercial longline fisheries were recorded. To standardize length across gears tunas measured in CFL were converted to FL using a regression equation developed by Farley et al. (2006). Fish ranged in size from 69.7 to 174.7 cm SFL. The minimum length in US waters for bigeye is 27 in. (68.58 cm) curved fork length, resulting in only large age 1 and 2 fish being retained. After extraction, sagittal otoliths were rinsed with water, dried, and stored in vials.

Otolith processing was based on methods developed by Busawon et al. (2015) and modified by Rodríguez-Marín (2019) for Atlantic bluefin tuna (*Thunnus thynnus*). All otoliths were cleaned in a jewelry sonicator to remove any remaining residual tissue that dried after extraction. Whole sagittal otoliths (left and right) were weighed and imaged, then embedded using Epothin 2 Epoxy hardener and resin at a 17:40 ratio respectively. Transverse sections were cut using an Isomet

1000 low-speed saw with diamond edged Buehler blades. Four transverse sections, 0.8 mm in width, were cut beginning with a rostral 'V-section' that included the origin.

Sections were mounted to glass slides using *QuickStick* Mounting Wax with the side closest to the origin facing down. After mounting, sections were polished using 180, 320 and 600 grit sandpaper to a width of roughly 0.3–0.5 mm. A final polish with a felt pad containing a light coating of water mixed with *MicroPolish 2* Alumina powder was applied to each section. Otolith sections were imaged under transmitted light with a compound microscope and features such as contrast and brightness were adjusted for each section in Adobe Photoshop. A 1 mm scale bar was created and placed at the first inflection point on the ventral arm of each bigeye section image to provide the reader guidance on the approximate location of first annulus formation based on mean distances in Farley et al. (2006).

All four sections were read twice by the same reader with no a priori information about the section (e.g., fish size, weight, previous age estimates). After analysis, only the two sections closest to the origin were used for age final estimation and reading error estimates. If age estimates from the first and second read were not identical, the section was aged again and assigned a final age based on that third reading. A readability score on a scale from 1 to 5 (1 = unreadable, 5 = excellent) was established and sections with low readability scores (mean score ≤ 2.5) were not included in final age estimates. Annual ages were assigned to individuals based on the number of fully formed opaque zones.

2.5.2. Otolith data from Hallier et al. (2005) (daily readings)

Data from a 2005 study published by Hallier included 255 bigeye tuna otoliths read for daily age (Hallier et al., 2005). The lengths ranged from 20 to 190 cm (FL) and the ages (in days) ranged from 116 to 3324. A light microscope was used for readings from fish with FL less than 74 cm and a scanning electron microscope for fish with FL equal to or bigger than 74 cm. The otolith preparation and reading protocols are described in detail in Hallier et al. (2005). Hallier's original study retained 83 of the 108 otoliths from Dakar, Senegal, and 147 from Abidjan, Ivory Coast. Given concerns regarding potential underaging of fish, in the current study the dataset was restricted to 153 age-length pairs from fish under 1 year old (Fig. 10). Fork lengths for this subset ranged from 29 to 67 cm.

2.5.3. Otolith data from the panama city lab

Twelve otoliths prepared and read by the Panama City Lab (PCL) were included in this study. Detailed protocols are described in Pacicco et al. (2021) and are aligned with the abovementioned annual ageing protocols (Allman et al., 2020). Ages determined by counting purported annual growth zones in these twelve Bigeye otolith cross sections were validated with bomb radiocarbon dating (Andrews et al., 2020). The valid age-at-length data for Bigeye were 3–17 years for fish lengths of 128.0–175.0 cm FL ($n = 12$) (Fig. 10).

2.6. Growth curve analyses

We fitted two different growth models to the tag-recapture data and counts of growth rings in otoliths: the Richards (1959) and the von Bertalanffy (1938) models. The Schnute (1981) parameterization was used as it allows for both models to be expressed using a single equation where the shape parameter, p , controls the level of curvature and reverts the function to a classic von Bertalanffy model when p is equal to 1.0.

All computations (except for the integrated analyses) were done using the R program language (R Core Team, 2020). Models based on just the otolith data were fitted using the R package 'FSA' (Ogle et al.,

2021). For the integrated analyses, we used the "Aires da Silva-Maunders-Schaefer-Fuller with correlation" (AMSFC) framework (Francis et al., 2016). This approach models the release and recapture lengths of fish as functions of age by treating age at tagging as a random effect. It also accounts for correlation between the measurements of an individual through the parameter ρ , which models correlation as a simple decreasing function of time at liberty (Δt ; Francis et al., 2016). The objective function is the sum of the bivariate normal log-likelihood of the release and recapture lengths, the lognormal log-likelihood of the random effects and the log-likelihood of the otolith data. Computer code in AD Model Builder (Fournier et al., 2012) was used that was based on the Bluefin tuna work of Ailloud et al. (2017).

For the Schnute (1981) model, the following equations are used:

$$L_a = \left((L_1)^p + ((L_2)^p - (L_1)^p) \frac{(1 - \exp(-K(a - A_1)))}{(1 - \exp(-K(A_2 - A_1)))} \right)^{\frac{1}{p}} \quad (1)$$

$$L_{a+\Delta t} = \left((L_1)^p + ((L_2)^p - (L_1)^p) \frac{(1 - \exp(-K(a + \Delta t - A_1)))}{(1 - \exp(-K(A_2 - A_1)))} \right)^{\frac{1}{p}} \quad (2)$$

$$L_\infty = \left(\frac{\exp(KA_2)(L_2)^p - \exp(KA_1)(L_1)^p}{\exp(KA_2) - \exp(KA_1)} \right)^{\frac{1}{p}} \quad (3)$$

$$t_0 = A_1 + A_2 - \frac{1}{K} \ln \left(\frac{\exp(KA_2)(L_2)^p - \exp(KA_1)(L_1)^p}{(L_2)^p - (L_1)^p} \right) \quad (4)$$

$$t^* = A_1 + A_2 - \frac{1}{K} \ln \left(p \frac{\exp(KA_2)(L_2)^p - \exp(KA_1)(L_1)^p}{(L_2)^p - (L_1)^p} \right) \quad (5)$$

where:

a is age,

L_a is the expected length of a fish of age a , thus $L_{a+\Delta t}$ is the expected length of a fish tagged at age a and recaptured at age $a + \Delta t$,

L_∞ is the asymptotic length,

K is the growth coefficient,

t_0 is the theoretical age at size 0 in the von Bertalanffy growth model ($p = 1$),

t^* is age at which the inflection of the Richards growth curve occurs ($p \neq 0$),

A_1 is age of youngest fish in sample,

A_2 is age of oldest fish in sample,

L_1 is the expected length of fish age A_1 ,

L_2 is the expected length of fish age A_2 .

In fitting the model, there are three types of parameters: those fixed by the user (A_1 and A_2), those estimated when maximizing the likelihood (L_1 , L_2 , and the parameters defining the error structure (see below)), and those derived from the other parameters (L_∞ , t_0 , and t^* (see above) and a^* and b^* (see below)). The parameters of the error structure are:

k_p steepness of slope ($k_p > 0$) defining relationship between correlation coefficient (ρ) and time at liberty (higher value means the faster the correlation coefficient declines to zero),

ρ_0 correlation (ρ) between length at tagging and length at recovery when time at liberty is zero ($0 < \rho_0 < 1$, and note that $\rho = 1 - \frac{1 - \rho_0}{1 - \rho_0 e^{(-k_p \Delta t)}}$ where Δt is time at liberty),

σ_{L_1} variability in length for fish at age A_1 ,

σ_{L_2} variability in length for fish at age A_2 ,

$\mu_{\log A_{\text{tag}}}$ - mean for random effects for age at tagging (follows lognormal distribution),

$\sigma_{\log A_{\text{tag}}}$ - standard deviation for random effects for age at tagging

Table 4

Parameter estimates from fitting von Bertalanffy models to the otolith data and also the von Bertalanffy results from the integrated model applied to the otolith and tagging data. Also shown are the parameter estimates when age-1 and age-2 fish were removed from the PFL otolith data for both models. The 95% Bootstrap confidence interval is given in parentheses for the parameter estimates from the otolith data only model. The standard deviation is given in parenthesis for the parameter estimates from the otolith and tagging data.

Parameter	Otolith Data only		Integrated Model, Otolith data + Tagging	
	All otoliths	PFL age 1 and 2 otoliths removed	All otoliths and tagging data	PFL age 1 and 2 otoliths removed and tagging data
K	0.464 (0.403, 0.543) 154.148 (147.081, 161.491) -0.163 (-0.250, -0.085)	0.392 (0.355, 0.441) 161.206 (154.389, 166.835) -0.239 (-0.306, -0.175)	0.271 (0.015) 178.700 (5.906) -0.537 (0.028)	0.252 (0.014) 185.780 (6.298) -0.524 (0.025)
L_{∞}				
t_0				

(follows lognormal distribution),

The derived parameters are:

a^* intercept for true variability around mean curve (variability in length at age) - linear model,

b^* slope for true variability around mean curve (variability in length at age) - linear model, (note - $\sigma_{L_a} = a^* + b^*$).

When $p = 1$, the model reverts to a von Bertalanffy model. Otherwise, the model assumes a Richards form.

2.7. Measurement error

It is possible to estimate the measurement error from short-term recaptures for fish with measured or estimated lengths (Ailloud et al., 2014). Define an increment, I , to be the length at the time of recapture, L_r , minus the length at the time of tagging, L_t . Over a suitably short time at liberty, the expected value of an increment is zero. We assume growth for any fish at liberty for less than Δ days is zero, the two recorded lengths are determined independently, the measurement error is the same at the time of tagging and recapture, and it does not vary with the length of the fish. Then the variance of the increments is

$$Var(I) = Var(L_r - L_t) = Var(L_r) + Var(L_t) = 2\sigma^2 \quad (6)$$

where $Var(L_r)$ and $Var(L_t)$ refer to the variance of repeated measurements of the same fish and is the measurement error. Hence, the measurement error standard deviation can be estimated by dividing the increment standard deviation by the square root of 2. If Δ is a short period of time, there is high assurance that growth while at liberty is close to zero at the cost of a smaller sample size compared to using a larger Δ . We Use $\Delta = 25, 50, 75$ and 98 days.

3. Results

3.1. Length measurement error

The Hallier et al. (2005) data have the lowest measurement error for length (4.7 or 4.9 cm depending on whether the cutoff Δ is set to 25 or 50 days) based on more than 500 paired measurements. The AOTTP data have slightly higher but similar measurement error (6.8 cm for both 25- and 50-day cutoffs, based on more than 900 paired measurements). The ICCAT database is extensive but there are very few short-term recaptures. With a cutoff Δ of 50 days, there are only 5 measurement pairs. With the 98-day cutoff, there are 26 pairs and the estimated measurement error is 10.9 cm. However, the mean size of the tagged fish increased by about 2 cm while at liberty so some of the estimated

measurement error might be due to unaccounted growth. With such a small sample size, this estimate is sensitive to outliers and the removal of a single datapoint reduces the estimated measurement error to 3.9 cm.

3.2. Growth curves fitted to otolith data

The nonlinear least squares estimates for models fitted to just otolith data are given in Table 4 along with non-parametric bootstrap estimates from 999 resampled datasets using the package 'nlstools' (Baty et al., 2015). Goodness of fit of the von Bertalanffy growth model can be judged from Fig. 10. The otoliths for the six oldest fish, four from the PCL data and two from the PFL data, are above the fitted growth curve when age-1 and age-2 fish from the PFL data are included in the study (red line). A sensitivity analysis was conducted on the influence of age-1 and age-2 fish from the PFL dataset. Those age-1 and age-2 fish had larger lengths than other age-1 and age-2 fish, indicating a potential sampling bias. All of the age-1 and age-2 fish from the PFL data were removed from the analysis ($n = 41$). Once these data points were removed the von Bertalanffy model was refit to the length-age pair data (Table 4, Fig. 10). The removal of the large age-1 and age-2 fish caused the estimate of asymptotic length to increase, the estimate of the time-axis intercept to decrease, and the estimate of K to decrease. When the larger age-1 and age-2 fish are removed, five of the oldest fish fall above the line and one below (Fig. 10, black line). Because the number of old fish in the dataset is limited, we fit the von Bertalanffy model while fixing the value of L_{∞} (between 145 and 200) to provide plausible pairs of K and L_{∞} for use in population models (see Table E1 of Appendix 5).

3.3. Integrated Analysis Results

The dataset consists of 1592 tag-recovery pairs from the AOTTP database, 18 tag-recovery pairs from ICCAT data, 146 tag-recovery pairs from Hallier dataset, 63 length-age pairs from AOTTP otoliths, 153 length-age pairs from otoliths from the Hallier dataset, 229 length-age pairs from PFL otoliths, and 12 length-age pairs from the PCL otoliths. Complete results with estimates for all of the parameters (fixed, estimated, and derived) can be found in (Table C1 of Appendix 3). The results from the Richards (Schnute with $p < 1$) and Von Bertalanffy (Schnute with $p = 1$) models were identical, and the Richards model estimated $p = 1.000$ (Table C1 of Appendix 3).

A sensitivity analysis was conducted on the influence of the 41 large age-1 and age-2 fish from the PFL dataset. The integrated model was fitted to the otolith data minus the 41 PFL fish (Table 4, Fig. 10, and Table C2 of Appendix 3). The integrated models, with or without the deletion of PFL's age 1 and age 2 fish, had higher asymptotic sizes than the corresponding models based just on the otolith data. All six of the oldest fish were below the integrated model curves. Similar to the results from the otolith only data, the removal of the large age-1 and age-2 fish from the integrated model caused the estimate of asymptotic length to increase and the estimate of K to decrease. However, in the integrated model, the estimate of the time-axis intercept became less negative when the age-1 and age-2 PFL fish were removed (Table 4).

Goodness of fit of the von Bertalanffy growth model from the integrated analysis with PFL age-1 and age-2 fish removed can be observed in Fig. 10 (green line) and Fig. 11. The integrated model describes the growth of young bigeye well but the growth of older fish tends to be above the predicted line (Fig. 11b). Otoliths for the six oldest fish, four from the PCL data and two from the PFL data, are below the fitted growth curves with and without the 1 and 2 year olds from PFL's data (Fig. 10). The vast majority of the data (nearly 100% of tagging and ~95% of length-age pairs) come from fish age-5 or younger.

The von Bertalanffy curve from just the otolith data (with age-1 and age-2 fish from PFL's otolith data removed) was plotted with the tagging data (Fig. 11a). This model fits the fish tagged at an older age better than the integrated model, but overestimates the growth for fish tagged at a young age with short times at liberty.




Fig. 11. (a). Vector plot of the tagging data from AOTTP, Hallier et al. (2005) and ICCAT. The vector plot is made by computing the predicted age for the length at tagging using the estimated von Bertalanffy growth parameters, and then assuming the age at recapture is the predicted age at tagging plus the time at liberty. The von Bertalanffy curve is based on the growth model from the otolith data only without the data for age-1 and age-2 fish from the PFL dataset. The fastest growing 1% and the slowest growing 1% (in cm/wk) of the records have been eliminated from the AOTTP data. One obvious outlier is seen among the 18 records from ICCAT. (b). Vector plot of the tagging data from AOTTP, Hallier et al. (2005) and ICCAT. The vector plot is made by computing the predicted age for the length at tagging using the estimated von Bertalanffy growth parameters, and then assuming the age at recapture is the predicted age at tagging plus the time at liberty. The von Bertalanffy curve is based on the integrated model applied to the data without age-1 and age-2 fish from the PFL otolith dataset. The fastest growing 1% and the slowest growing 1% (in cm/wk) of the records have been eliminated from the AOTTP tagging data. One obvious outlier is seen among the 18 records from ICCAT tagging data; this record was not used to fit the von Bertalanffy model.

4. Discussion

By utilizing data from both tagging studies and length-age pairs we were able to estimate several models for growth of bigeye tuna in the Atlantic Ocean. We incorporated multiple datasets to estimate a comprehensive growth model for the species.

The estimates of length measurement error in the AOTTP tagging data (roughly 7 cm) are similar to those found by Ailloud et al. (2014) for Bluefin tuna, i.e., roughly 5 cm. Measurement error in the tagging study of Hallier et al. (2005) was about 4.8 cm, slightly better than that in the AOTTP bigeye tuna data and the Bluefin tuna data. Unfortunately, the ICCAT tagging database for bigeye tuna contains very few short-term recaptures making it difficult to assess measurement error. Visual examination of the ICCAT tagging data (Fig. 11) showed one obvious outlier and slightly slower growth than the tagging data of Hallier et al. (2005) and the ICCAT tagging data. There was no justification why these 17 records (after removal of the outlier) were invalid so they were retained in the analysis.

The integrated model runs yielded similar results between Richards and Von Bertalanffy when using the same datasets, i.e., the estimated value of the shape parameter p was close to or equal to the value of 1.0 at which the Schnute model reverts to a von Bertalanffy curve. The integrated model run using the von Bertalanffy model and all of the data (Table 4) estimates an L_{∞} of 178.70 cm FL (SD 5.906) and K of 0.271 yr^{-1} (SD 0.015). The integrated model run removing PFL's age-1 and age-2 fish estimates an L_{∞} of 185.78 cm FL (SD 6.298) and a K of

0.252 yr^{-1} (SD 0.014). These results are similar to those found in previous studies (Fig. A1 of Appendix 1). It is worth noting that there was only one other integrated study completed for bigeye using tagging and otolith data (Hallier et al., 2005). The integrated results from that study yielded a larger estimate of L_{∞} (217.28 cm FL) and a smaller estimated K (0.180 yr^{-1}). One explanation is that Hallier used daily ring readings for the otoliths beyond 1 year, a practice which has been shown to be unreliable for bigeye (Williams et al., 2013; Krusic-Golub and Ailloud (n.d.)). The Hallier study also used a much shorter time at liberty cutoff (14 days) versus the 98 days used here, had few old fish, and few long-term recaptures. Our results are more similar to the SS3 fits to the Hallier et al., 2005 data used by the 2018 ICCAT bigeye stock assessment (Anon, 2019) $L_{\infty} = 179.9$ and $K = 0.281$.

Of particular note is that we were able to extend the maximum age used in the analysis up to 17 years compared to the maximum age of approximately 9 years used by Hallier et al. (2005) to estimate the growth parameters currently used in the stock assessment and the maximum age of 8 found in the study by Delgado de Molina and Santana (1986). The importance of this is amplified by the lack of tag returns from either very large fish or fish at liberty for a very long time, i.e., from old fish.

Normally, it would be prudent to analyze different data sources independently to identify any conflicts among data sources. However, when fitting an asymptotic growth model, it is essential to have a wide contrast in the independent variable, age. It is impossible to judge the rate of curvature with change in age if age does not change (much)

among observations. Lack of old animals in a growth study often leads to extremely high estimates of asymptotic size whereas lack of young animals can lead to very large, negative estimates of the t_0 parameter. In the case of bigeye tuna, the use of multiple datasets captured a broad size spectrum of the population but the use of just the tagging data resulted in very few large, old fish.

The study of Hallier et al. (2005) used otoliths and tagging data while the study of Delgado de Molina and Santana (1986) used growth rings in dorsal spines. The latter authors noted problems with remodeling of the central cavity of the spine which resulted in the loss of rings representing the first years of life. Given the current study utilized several datasets over more than two decades, contained the widest range of sizes, and included validated ages beyond those currently assumed by ICCAT, this new curve should represent the most realistic estimates of bigeye tuna growth in the Atlantic to date.

The new curve with tagging and otolith data is similar to the curve used in the last stock assessment but with a lower estimated asymptotic size (Fig. 1). The six oldest fish in the study are below the estimated growth curve which suggests that the addition of additional old fish (> 7 yr, implying an effort to sample fish > 150 cm FL) to the analysis might bring the asymptotic size down and increase the growth coefficient K estimate.

Additional old fish should be collected (both from tagging and length-age data) in order to better estimate the model. The curve based on otolith data (without PFL's age 1 and age 2 fish) goes through the cloud of six old fish on the right. In the absence of adequate samples of old (large) fish, one can artificially assign greater weight to the existing samples of old fish to force the curve to go through the cloud of data points of old fish (e.g., Maunder et al., 2018). We estimated the growth parameters K and t_0 with L_∞ held fixed at various values (Appendix 5). This shows that the asymptotic size is not well determined, with fits having L_∞ fixed anywhere between 155 and 170 cm FL having similar residual standard errors when age 1 and age 2 PFL fish are removed from the dataset.

All of the tagging data appear to be in some conflict with the otolith data (Figs. 11a, 11b). When the vector plot is made with the curve fitted to just otolith data (Fig. 11a), the young fish appear to grow slower than predicted by the growth curve as evidenced by the observation that there are more termini of the vectors to the right (below) the fitted line than to the left (above); fish recaptured at an older age (age > 2) tend to be to the left (above) the fitted line. This suggests a conflict between the tagging and otolith results. When the curve is fitted to tagging and otolith data, a different pattern appears in the vector plot (Fig. 11b). Now, the vectors for fish tagged at age 0 and age 1 appear to be symmetric about the regression line, but the lack of fit for fish recaptured at age > 2 is worse than in Fig. 11a.

It is not clear why the tagging and otolith data are in disagreement. We propose the model based on otolith data provides the most realistic estimates of bigeye tuna growth because it predicts the size of old fish through the fitted value of L_∞ and it avoids patterns in the residuals from the tagging data (Appendix 4). If tagged bigeye tuna with longer times at

liberty are recaptured in the future it could resolve the apparent discrepancy between the tagging and otolith data. The inclusion of additional otoliths and tag returns from old fish would improve both models as the sample size for old fish remains limited.

CRedit authorship contribution statement

Lynn Waterhouse: Conceptualization, Methodology, Writing – original draft, Writing – review & editing, Formal Analysis, Funding acquisition, Visualization. **Lisa Ailloud:** Writing – original draft, Writing – review & editing, Resources, Data curation. **Riley Austin:** Writing – review & editing, Resources, Investigation. **Walter J. Golet:** Writing – review & editing, Resources, Investigation. **Ashley Pacicco:** Resources, Investigation. **Allen H. Andrews:** Writing – review & editing, Resources, Investigation. **Khady Diouf:** Resources, Investigation. **Yacine Ndiour:** Resources, Investigation. **Kyne Krusic-Golub:** Resources, Investigation. **Guelson da Silva:** Writing – review & editing, Resources, Investigation. **John M. Hoenig:** Conceptualization, Methodology, Writing – original draft, Writing – review & editing, Funding acquisition, Visualization.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank members of the AOTTP tag and recovery teams, including biological samplers, for doing the meticulous work of otolith extraction in recovered fish. We thank Captains Billy McIntyre of the F/V Shady Lady, Dan Mears Jr. of the F/V Monica, and their respective crew who were vital for longline sampling efforts and personal communications regarding fishery dynamics. Officials and participants in the Oak Bluffs Bluewater Classic, Mid-Atlantic Tuna, Ocean City Tuna, White Marlin Open, and Mid-Atlantic Billfish tournaments were also crucial for obtaining samples. Graduate students and lab technicians, Kelsey Moon, Isabelle See, Samantha Nadeau, and Brenda Rudnicki assisted with field and lab work for this project as well. Funding to support biological sampling of tunas, age estimation and radiocarbon analysis was provided by NOAA CRP # NA17NMF4540140 to Walt Golet in the School of Marine Sciences at the University of Maine. This work was also supported by the European Union (DCI-FOOD/2015/361-161). Additional financial support from ICCAT Contracting Parties and Cooperating non-Contracting Parties is gratefully acknowledged. Funding provided for L. Waterhouse and J. Hoenig by International Commission for the Conservation of Atlantic Tunas under fiscal identification number N4001546C. The anonymous reviewers and Beth Matta provided helpful comments.

Appendix 1

See Appendix Table A1.

Table A1Estimates of growth parameters for bigeye tuna in the Atlantic Ocean. L_{∞} and Length Range pertain to fork length in cm; K is in units of yr⁻¹ and t_0 is in yr.

Growth Function	L_{∞}	K	t_0	Length Range	Method	Reference
VB	161.21	0.392	-0.239	29 – 175	Otoliths	This study
VB	185.78	0.252	-0.524	28–175	Otoliths, tagging	This study
Peterson's method	338.53	0.10497	-0.5425	58–140	Length Frequency (predorsal)	Champagnat and Pianet (1974)
VB	199.77	0.1970	-0.71		Spines, males (n = 245)	Delgado de Molina and Santana (1986)
VB	214.54	0.1669	-0.77		Spines, females (n = 239)	Delgado de Molina and Santana (1986)
VB	206.14	0.1822	-0.74	50–180	Spines, both sexes (n = 540)	Delgado de Molina and Santana (1986)
VB	253.75	0.173	-0.15		Spines	Gaikov et al. (1980)
VB	491.6	0.0135	3.808	40–180	Length Frequency	Weber (1980)
VB	218.8	0.23	-0.20	56–190	Spines (n = 77)	Draganik and Peiczarski (1984)
VB	195.54	0.206		37 – 124	Tagging (n = 625)	Hallier et al. (2005)
VB (solved in excel)	207.43	0.202	-0.613	29 – 190	Otoliths, daily (n = 230)	Hallier et al. (2005)
VB (FAO vonbit)	206.976	0.203	-0.616	29 – 190	Otoliths, daily (n = 230)	Hallier et al. (2005)
Gompertz	179.13	0.4088	(A= 1.7268) (b= -7.185, m= 2280.4)	29 – 190	Otoliths, daily (n = 230)	Hallier et al. (2005)
Richards	178.63	0.424		29 – 190	Otoliths, daily (n = 230)	Hallier et al. (2005)
VB	217.28	0.180	-0.709	37 – 124 & 29 – 190	Tagging (n = 625) & Otoliths, daily (n = 230). Used in last stock assessment.	Hallier et al. (2005)
VB	264.02	0.12	-0.68	44 – 179	Caudal vertebrae (n = 175)	Alves et al. (2002)
VB	285.3745	0.1127			Tagging (n = 243)	Cayré and Diouf (1984)
VB / Petersen's method	249.6	0.0124	-4.78		Length frequency	Marcille et al. (1978)

Appendix 2

See Appendix [Tables B1 and B2](#) here.

Table B1Definition of length measurements used in conversions for tuna lengths ([Table B2](#) of Appendix 2).

Length	Type	Description
CFL	Curved fork length	
CKL	Cleithrum-Keel length	Projected straight distance between the point on the cleithrum that provides the shortest possible measurement to the anterior portion of the caudal keel. The cleithrum is the semicircular bony structure at the posterior edge of the gill opening.
EYF	Eye to fork	
FL	Fork Length	Projected straight distance from the tip of the upper jaw (snout) to the shortest caudal ray (fork)
FLT	Curved Fork Length	Projected curved-body distance from the tip of the upper jaw (snout) to the shortest caudal ray (fork).
LD1	Lower jaw to 1st dorsal	
LJF	Lower Jaw Fork Length	* **Equivalent to fork length for tropical tuna species.
PFL	Pectoral-fork length	Projected straight distance between the most anterior insertion of the pectoral fin and the fork of the tail.
PFLT	Curved Pectoral-fork length	
SFLT	Straight fork length	Projected curved-body distance between the most anterior insertion of the pectoral fin and the fork of the tail.
SL	Snout (preorbital) length	* **Equivalent to fork length for tropical tuna species.
STD	Standard length	
TLE	Total length	
UNK	Unknown	

Source: Modified from [Table 1](#) in [IOTC Secretariat \(2006\)](#) and [Scida et al. \(2001\)](#).

Table B2

Length conversions used to convert between different length standards. Acronyms and definitions are given in [Table B1](#) of Appendix 2.

L	a	b	R2	Standard Length (Y = a+b*L)	Area	Range (cm)	n	Ref
PFL	18.191	1.2129	0.8988	FL	Atl.	33–141	3174	1
CKL	-5.5109	0.6215	0.9255	FL	Atl.	29–110	570	1
PFLT	-2.287	1.4572	0.9564	FL	Atl.	44–110	59	1
PFLT	7.1818	1.3418	0.9733	FLT	Atl.	44–110	59	1
FLT	0.9082	0.9676	0.9891	FL	Atl.	63–169	304	1

Appendix 3

Complete results from fitting the integrated model to all of the data ([Table C1](#) Appendix 3) and to the data with age-1 and age-2 fish from PFL data removed ([Table C2](#) Appendix 3).

Table C1

Results from Richards and Von Bertalanffy models fitted to the full data set consisting of tagging data from AOTTP, ICCAT, and the [Hallier et al. \(2005\)](#) study plus hard part data from AOTTP, Hallier (only ages < 1 yr), PFL data, and PCL data. Symbols are defined in the main text. Note that p is fixed at 1.0 in the von Bertalanffy model and estimated to be 1.000 in the Richards model.

	Richards		Von Bertalanffy	
	(Schnute with $ p < 100$)		(Schnute with $p = 1$)	
	Value	S.E.	Value	S.E.
Fixed parameters				
A_1	0	–	0	–
A_2	17	–	17	–
p	–	–	1	–
Estimated parameters				
L_1	24.364	0.765	24.364	0.765
L_2	178.910	5.560	178.910	5.560
K	0.266	0.015	0.266	0.015
p	1.00	8.867e-08	–	–
k_p	1.381	0.232	1.381	0.232
ρ_0	0.887	0.015	0.887	0.015
σ_{L_1}	2.220	0.387	2.220	0.387
σ_{L_2}	26.713	1.063	26.713	1.063
$\mu_{\log \text{Atag}}$	-0.187	0.020	-0.187	0.020
$\sigma_{\log \text{Atag}}$	-1.068	0.034	-1.068	0.034
Derived parameters				
L_∞	180.590	6.046	180.590	6.046
t_0	–	–	-0.544	0.028
t^*	-0.544	0.028	–	–
a^*	-1.641	0.500	-1.641	0.500
b^*	0.158	7.711e-03	0.158	7.711e-03
b	-1.000	7.679e-08	–	–
Negative log-likelihood	9238.97945		9238.97945	

Table C2

Results from Richards and Von Bertalanffy models fitted to the full data set consisting of tagging data from AOTTP, ICCAT, and the Hallier et al. (2005) study plus hard part data from AOTTP, Hallier (only ages < 1 yr), PFL data, and PCL data. Symbols are defined in the main text. The age-1 and age-2 fish have been removed from PFL ($n = 41$) to avoid probable sampling bias. Note that p is fixed at 1.0 in the von Bertalanffy model and estimated to be 1.000 in the Richards model.

	Richards		Von Bertalanffy	
	(Schnute with $ p < 100$)		(Schnute with $p = 1$)	
	Value	S.E.	Value	S.E.
Fixed parameters				
A_1	0	–	0	–
A_2	17	–	17	–
p	–	–	1	–
Estimated parameters				
L_1	23.127	0.642	23.127	0.642
L_2	185.450	5.790	185.450	5.790
K	0.247	0.014	0.247	0.014
p	1.000	1.317e-07	–	–
k_p	1.434	0.253	1.434	0.253
ρ_0	0.864	0.019	0.864	0.019
σ_{L_1}	1.571	0.322	1.571	0.322
σ_{L_2}	26.297	1.074	26.297	1.074
$\mu_{\log \text{Atag}}$	-0.132	0.017	-0.132	0.017
$\sigma_{\log \text{Atag}}$	-1.076	0.031	-1.076	0.031
Derived parameters				
L_∞	187.900	6.455	187.900	6.455
t_0	–	–	-0.531	0.025
t^*	-0.531	0.025	–	–
a^*	-1.952	0.429	-1.952	0.429
b^*	0.152	7.055e-03	0.152	7.055e-03
b	-1.000	1.178e-07	–	–
Negative log-likelihood	9014.8878		9014.8878	

Appendix 4

Residual plots from the preferred model (length-age pair data only) and the best fitting integrated model. For the preferred model (length-age pair data only), the asymptotic length L_∞ (fork length) equals 161.21 cm (95% bootstrap CI 154.39, 166.84), growth parameter K equals 0.392 yr^{-1} (95% bootstrap CI 0.355, 0.441), and the time-axis intercept t_0 equals -0.239 yr (95% bootstrap CI -0.306 , -0.175). For the best fitting integrated model, the asymptotic length L_∞ (fork length, in cm) was estimated to be 185.78 (SD 6.298), the growth parameter K was 0.252 yr^{-1} (SD 0.014), and the time-axis intercept t_0 was -0.524 yr (SE 0.025). Figs. A1–A4.




Fig. A1. Residual plot by age from the preferred model (length-age pair data only).




Fig. A2. Residual plot by age from the integrated model for the length-age pair data only.




Fig. A3. Residual plot by recovery fork length (cm) from the integrated model for the tagging data only. A release age was calculated using the parameter estimates and the release fork length (cm). The recovery age was calculated by adding time at liberty to the calculated release age.




Fig. A4. Residual plot by estimated recovery age from the integrated model for the tagging data only. A release age was calculated using the parameter estimates and the release fork length (cm). The recovery age was calculated by adding time at liberty to the calculated release age.

Appendix 5

Parameter estimates for the von Bertalanffy growth model (length-age pair data only) were computed when the value of the asymptotic length L_{∞} (fork length) was fixed. When the model is fit without setting a value for asymptotic length L_{∞} and with PFL age 1 and 2 otoliths removed (Table 4), the asymptotic length L_{∞} (fork length) equals 161.21 cm (95% bootstrap CI 154.39, 166.84), growth parameter K equals 0.392 yr^{-1} (95% bootstrap CI 0.355, 0.441), and the time-axis intercept t_0 equals -0.239 yr (95% bootstrap CI -0.306 , -0.175). From Fig. 10 the values of the asymptotic length L_{∞} (fork length) that would best fit the oldest fish would be between 165 and 175, and this would result in a value of growth parameter K between 0.368 and 0.316 (when PFL age 1 and 2 otoliths are removed) or 0.380–0.325 (when all otoliths are included). Fig. E1 and Table E1.




Fig. E1. Estimated value of von Bertalanffy growth parameter K , and residual standard error, when the curve is fit to the data without the PFL age 1 and age 2 fish and with the value of L_{∞} fixed.

Table E1

Parameter estimates from fitting von Bertalanffy models to the otolith data when fixing the value of asymptotic length L_{∞} (fork length). This was done with otolith data when the age-1 and age-2 fish were removed from the PFL otolith data and with all otolith data. Estimated values for growth parameter K and time-axis intercept t_0 and the residual standard error have been rounded to three decimal places.

Set Value L_{∞}	PFL age 1 and 2 otoliths removed			All otoliths		
	Estimated Value			Estimated Value		
	K	t_0	Residual Standard Error	K	t_0	Residual Standard Error
145	0.540	-0.078	9.292	0.564	-0.064	10.344
150	0.484	-0.130	8.987	0.505	-0.118	10.202
155	0.439	-0.180	8.817	0.456	-0.172	10.170
160	0.400	-0.228	8.752	0.415	-0.224	10.218
165	0.368	-0.274	8.769	0.380	-0.275	10.324
170	0.340	-0.319	8.850	0.351	-0.324	10.472
175	0.316	-0.362	8.980	0.325	-0.371	10.650
180	0.294	-0.403	9.147	0.302	-0.417	10.849
185	0.276	-0.443	9.341	0.283	-0.461	11.063
190	0.259	-0.481	9.555	0.265	-0.504	11.286
195	0.244	-0.519	9.782	0.249	-0.546	11.514
200	0.231	-0.555	10.018	0.235	-0.586	11.744

References

- Ailloud, L.E., Lauretta, M.V., Hoenig, J.M., Walter, J.F., Fonteneau, A., 2014. Growth of atlantic bluefin tuna determined from the ICCAT tagging database: a reconsideration of methods. Collect. Vol. Sci. Pap. ICCAT 70, 380–393.
- Ailloud, L.E., Lauretta, M.V., Hanke, A.R., Golet, W.J., Allman, R., Siskey, M.R., Secor, D. H., Hoenig, J.M., 2017. Improving growth estimates for western Atlantic Bluefin tuna using an integrated modeling approach. Fish. Res. 191, 17–24.
- Ailloud, L.E., Beare, D., Farley, J.H. and Krusic-Golub, K., 2019. Preliminary results on AOTTP validation of otolith increment deposition rates in yellowfin tuna in the Atlantic. Collective Volume of Scientific Papers. ICCAT. 76(6):156–163.
- Aires-da-Silva, A.M., Maunder, M.N., Schaefer, K.M., Fuller, D.W., 2015. Improved growth estimates from integrated analysis of direct aging and tag-recapture data: an illustration with bigeye tuna (*Thunnus obesus*) of the eastern Pacific Ocean with implications for management. Fish. Res. 163, 119–126.
- Allman, R., Ailloud, L., Austin, R., Falterman, B., Farley, J., Lang, E., Pacicco, A., & Satoh, K. (2020). Report of the International Workshop On the Ageing of Yellowfin and Bigeye Tuna. Collective Volume of Scientific Papers. ICCAT. 77(8), 32–46.
- Alves, A., de Barros, P., Pinho, M.R., 2002. Age and growth studies of bigeye tuna *Thunnus obesus* from Madeira using vertebrae. Fish. Res. 54, 389–393.
- Andrews, A.H., Pacicco, A., Allman, R., Falterman, B.J., Lang, E.T., Golet, W., 2020. Age validation of yellowfin (*Thunnus albacares*) and bigeye (*Thunnus obesus*) tuna of the northwestern atlantic ocean. Can. J. Fish. Aquat. Sci. 77, 637–643.
- Anon. 2019. Report of the 2018 ICCAT Bigeye Tuna Stock Assessment Meeting. Collective Volume of Scientific Papers. ICCAT. 75(7): 1721–1855.
- Anon. 2021. ICCAT Atlantic Ocean tropical Tuna Tagging Programme (AOTTP) evidence based approach for sustainable management of tuna resources in the Atlantic. Final Report. Available at: (<https://www.iccat.int/aottp/AOTTP-Documents-Library/Reports/Interim-Report/AOTTP-Interim-Report-2019.pdf>).
- Baty, F., Ritz, C., Charles, S., Brutsche, M., Flandrois, J.-P., Delignette-Muller, M.-L., 2015. A toolbox for nonlinear regression in R: the package nlstools. J. Stat. Softw. 66, 1–21.
- Beare, D., Ailloud, L.E., Garcia, J., Pastor, R., Kebe, S., 2019. ICCAT Atlantic Ocean tropical Tuna Tagging Programme (AOTTP): Evidence based approach for sustainable management of tuna resources in the Atlantic. Available at: (<https://www.iccat.int/aottp/AOTTP-Documents-Library/Reports/Interim-Report/AOTTP-Interim-Report-2019.pdf>).

- Busawon E. D.S., Rodríguez-Marin P.L., Luque R., Allman B., Gahagan W., Golet E., Koob M., Siskey M.R., Sobrón P., Quelle J., Nielson D.H. Secor Evaluation of an Atlantic bluefin tuna otolith reference collection Collect. Vol. Sci. Pap. ICCAT 71 2 2015 960–982.
- Cayré, P., Diouf, T., 1984. Croissance du thon obèse (*Thunnus obesus*) de l'atlantique d'après les résultats de marquage. Collective Volume of Scientific Papers. ICCAT. 20: 180–187.
- Cayré, P., Amon Kothias, J.B., Diouf, T., Stretta, J.M., 1993. Biology of tuna. Pages 147–244 in A. Fonteneau and J. Marcille, editors. Resources, fishing and biology of the tropical tunas of the Eastern Central Atlantic. FAO Fisher. FAO, Rome.
- Champagnat, C., and R. Pianet. 1974. Croissance du Fatudo (*Thunnus obesus*) Dans Les Recions de Dakar et de Pointe-Noire Collective Volume of Scientific Papers. ICCAT. 2:141–144.
- Collette, B.B. C.E. Nauen. 1983. FAO species catalogue. Scombrids of the world. An annotated and illustrated catalogue of tunas mackerels, bonitos and related species know to date.
- Draganik, B., and W. Peiczarski. 1984. Growth and Age of Bigeye and Yellowfin Tuna in the Central Atlantic as Per Data Gathered by R/V Wicczno. Collective Volume of Scientific Papers. ICCAT. 20:96–103.
- Eveson, J.P., Laslett, G.M., Polacheck, T., 2004. An integrated model for growth incorporating tag-recapture, length-frequency, and direct aging data. Can. J. Fish. Aquat. Sci. 61, 292–306.
- Farley, J.H., Williams, A.J., Clear, N.P., Davies, C.R., Nicol, S.J., 2013. Age estimation and validation for South Pacific Albacore *Thunnus alalunga*. Journal of Fish Biology 82, 1523–1544.
- Farley, J.H., Clear, N.P., Leroy, B., Davis, T.L.O., McPherson, G., 2006. Age, growth and preliminary estimates of maturity of bigeye tuna, *Thunnus obesus*, in the Australian region. Mar. Freshw. Res. 57 (7), 713–724. <https://doi.org/10.1071/MF05255>.
- Fournier, D.A., Skaug, H.J., Ancheta, J., Ianelli, J., Magnusson, A., Maunder, M.N., Nielsen, A., Sibert, J., 2012. AD Model Builder: using automatic differentiation for statistical inference of highly parameterized complex nonlinear models. Optim. Methods Softw. 27, 233–249.
- Francis, R.I.C.C., 1988a. Maximum likelihood estimation of growth and growth variability from tagging data. N.Z. J. Mar. Freshw. Res. 22 (1), 43–51.
- Francis, R.I.C.C., 1988b. Are growth parameters estimated from tagging and age-length data comparable? Can. J. Fish. Aquat. Sci. 45 (6), 936–942.
- Francis, R.I.C.C., Aires-da-Silva, A.M., Maunder, M.N., Schaefer, K.M., Fuller, D.W., 2016. Estimating fish growth for stock assessments using both age-length and tagging-increment data. Fish. Res. 180, 113–118.
- Gaikov, V.V., Chur, V.N., Zharov, V.L., Fedoseev, Y.P., 1980. On age and growth of atlantic bigeye tuna. Collect. Vol. Sci. Pap. ICCAT 9, 294–302.
- Hallier, J.-P., Stequert, B., Maury, O., Bard, F.-X. 2005. Growth of bigeye tuna (*Thunnus obesus*) in the eastern atlantic ocean from tagging-recapture data and otolith readings. Collective Volume of Scientific Papers. ICCAT. 57:181–194.
- ICCAT Secretariat. (n.d.). Access to ICCAT statistical databases. (<https://www.iccat.int/en/accessingdb.html>).
- IOTC Secretariat. 2006. Biological data on tuna and tuna-like species gathered. IOTC-2006-WPB-INF01. at the IOTC Secretariat: Status Report | IOTC.
- Krusic-Golub, K., L.E. Ailloud. (n.d.). Evaluating otolith increment deposition rates in Atlantic Ocean bigeye and yellowfin tuna tagged during the Atlantic Ocean tropical Tuna Tagging Program.
- Lang, E.T., Falterman, B.J., Kitchens, L.L., Marshall, C.D., 2017. Age and growth of Yellowfin Tuna (*Thunnus albacares*) in the northern Gulf of Mexico. Collect. Vol. Sci. Pap. ICCAT 73 (1), 423–433.
- Laslett, G.M., Eveson, J.P., Polacheck, T., 2002. A flexible maximum likelihood approach for fitting growth curves to tag recapture data. Can. J. Fish. Aquat. Sci. 59 (6), 976–986.
- Lehodey P., Leroy B (1999) Age and growth of yellowfin tuna (*Thunnus albacares*) from the western and central Pacific Ocean as indicated by daily growth increments and tagging data. In: Proceedings 12th Standing committee on tuna and billfish working, pp1–21.
- Delgado de Molina, A., Santana, J.C., 1986. Estimacion de la edad y crecimiento del patudo (*Thunnus obesus*, Lowe 1939) capturado en las islas canarias. Collect. Vol. Sci. Pap. ICCAT 25, 130–137.
- Marcille, J., Champagnat, C., Armada, N., 1978. Croissance du patudo (*Thunnus obesus*) de l'Océan Atlantique intertropical oriental. Doc. Sci. Cent. De Rech. Océanographiques, Abidj. 9, 73–81.
- Maunder, M.N., Deriso, R.B., Schaefer, K.M., Fuller, D.W., Aires-da-Silva, A.M., Mente-Vera, C.V., Campana, S.E., 2018. The growth cessation model: a growth model for species showing a near cessation in growth with application to bigeye tuna (*Thunnus obesus*). Mar. Biol. 165 (76) <https://doi.org/10.1007/s00227-018-3336-9>.
- Ogle DH, Doll JC, Wheeler P., Dinno A. (2021). FSA: Fisheries Stock Analysis. R package version 0.9.1, (<https://github.com/droglenc/FSA>).
- Pacicco, A.E., Allman, R.J., Lang, E.T., Murie, D.J., Falterman, B.J., Ahrens, R. and Walter, J.F., III (2021), Age and Growth of Yellowfin Tuna in the U.S. Gulf of Mexico and Western Atlantic. Marine and Coastal Fisheries, 13: 345 - 361. <https://doi.org/10.1002/mcf2.10158>.
- R Core Team, 2020. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Richards, F.J., 1959. A flexible growth function for empirical use. J. Exp. Bot. 10, 290–301.
- Rodríguez-Marin, E., Landa, J., Ruiz, M., Godoy, D., Rodríguez- Cabello, C. (2004). Age estimation of adult bluefin tuna (*Thunnus thynnus*) from dorsal spine reading. Collective Volume of Scientific Papers. ICCAT. 56:1168–1174.
- Sardenne F et al (2015) Determining the age of tropical tunas in the Indian ocean from otolith microstructures. Fish Res 163:44–57. <https://doi.org/10.1016/j.fishres.2014.03.008>
- Schnute, J., 1981. A versatile growth model with statistically stable parameters. Can. J. Fish. Aquat. Sci. 38 (9), 1128–1140.
- Scida, P., A. Rainosek, and T. Lowery. 2001. Length conversions for yellowfin tuna (*Thunnus albacares*) caught in the western north atlantic ocean. Collective Volume of Scientific Papers. ICCAT. 52:528–532.
- Shuford, R.L., Dean, J.M., Stéquert, B., Morize, E., 2007. Age and growth of yellowfin tuna in the Atlantic Ocean. ICCAT Collective Volume of Scientific Papers 60, 330–341.
- Von Bertalanffy, L., 1938. A quantitative theory of organic growth (inquiries on growth laws. II). Hum. Biol. 10, 181–213.
- Weber, E. 1980. An Analysis of Atlantic Bigeye Tuna (*Thunnus obesus*) Growth. Collective Volume of Scientific Papers. ICCAT. 9:303–307.
- Williams, A.J., Leroy, B.M., Nicol, S.J., Farley, J.H., Clear, N.P., Krusic-Golub, K., Davies, C.R., 2013. Comparison of daily- and annual- increment counts in otoliths of bigeye (*Thunnus obesus*), yellowfin (*T. albacares*), southern bluefin (*T. maccoyii*) and albacore (*T. alalunga*) tuna. ICES J. Mar. Sci. 70, 1439–1450.