

**State of California
The Resources Agency
DEPARTMENT OF FISH AND GAME**

2011 CDFG FRGP REPORT

**LOWER REDWOOD CREEK
JUVENILE SALMONID (SMOLT) ABUNDANCE PROJECT
2004 – 2011 Seasons
CDFG PROJECT 2a7**

Fisheries Restoration Grants Program (Project Numbers: P0810509, S02067)

FINAL

Prepared by

**Michael D. Sparkman
Northern Region**

Anadromous Fisheries Resource Assessment and Monitoring Program

January 30, 2012

TABLE OF CONTENTS

Section	Page
LIST OF FIGURES	iii
LIST OF TABLES	iv
LIST OF APPENDICES.....	vi
ABSTRACT.....	1
INTRODUCTION	2
Site Description.....	3
Purpose.....	7
METHODS AND MATERIALS.....	8
Trap Operations	8
Biometric Data Collection	9
Population Estimates.....	10
Physical Data Collection.....	10
Statistical Analyses	11
RESULTS	12
Species Captured.....	12
Trapping Efficiencies.....	16
Population Estimates.....	17
Age Composition of Juvenile Steelhead Trout	33
Fork Lengths and Weights	34
Developmental Stages.....	40
Trapping Mortality.....	40
Stream Temperatures	42
DISCUSSION	45
0+ Chinook Salmon	45
1+ Chinook Salmon	48
0+ Steelhead Trout.....	49
1+ Steelhead Trout.....	51
2+ Steelhead Trout.....	54
Cutthroat Trout.....	56
0+ Coho Salmon	58
1+ Coho Salmon	59
0+ Pink Salmon.....	60
CONCLUSIONS.....	61
RECOMMENDATIONS.....	63
ACKNOWLEDGEMENTS	64
LITERATURE CITED	64
PERSONAL COMMUNICATIONS.....	72
APPENDICES	73

LIST OF FIGURES

Figure 1. Redwood Creek basin with rotary screw trap location (RM 4), Humboldt County, CA. (scale is slightly inaccurate due to reproduction process, Charlotte Peters pers. com. 2001).....	4
Figure 2. Total juvenile salmonid trap catches (n = 70,489) from April 9 th through September 9 th , 2011, lower Redwood Creek, Humboldt County, CA. Numeric values above columns represent actual catches. 0+ KS = young-of-year Chinook salmon, 1+ KS = age 1 Chinook salmon, 0+ SH = young-of-year steelhead trout, 1+ SH = age 1 and older steelhead trout, 2+ SH = age 2 and older steelhead trout, 0+ CO = young of year coho salmon, 1+ CO = age 1 and older coho salmon, CT = cutthroat trout, 0+ Pink = young-of-year pink salmon.	12
Figure 3. 0+ Chinook Salmon population abundance estimates (error bars are 95% confidence interval) in YRS 2004 – 2011. Numeric values next to box represent number of individuals. Line of best fit is a regression line with corresponding equation, correlation value (r), and p value, lower Redwood Creek, Humboldt County, CA.	17
Figure 4. Comparison of 0+ Chinook salmon population abundance by month in YR 2011 with the previous seven year average, lower Redwood Creek, Humboldt County, CA.	18
Figure 5. 0+ Chinook salmon population abundance by week in YR 2011, lower Redwood Creek, Humboldt County, CA.	19
Figure 6. 1+ Chinook Salmon population abundance estimates (error bars are 95% confidence interval) in YRS 2009 – 2011. Lack of 95% CI for YR 2011 is due to scale of ‘Y’ axis. Numeric values next to box represent number of individuals, lower Redwood Creek, Humboldt County, CA.	21
Figure 7. 1+ steelhead trout population abundance estimates (error bars are 95% confidence interval) in YRS 2004 – 2011. Numeric values next to box represent number of individuals. Line of best fit is a regression line, with corresponding equation, correlation value (r), and p value, lower Redwood Creek, Humboldt County, CA.	22
Figure 8. Comparison of 1+ steelhead trout population abundance by month in YR 2011 with the previous seven year average, lower Redwood Creek, Humboldt County, CA.	23
Figure 9. 2+ steelhead trout population abundance estimates (error bars are 95% confidence interval) in YRS 2004 – 2011. Numeric values next to box represent number of individuals. Line of best fit is a regression line, with corresponding equation, correlation value (r), and p value, lower Redwood Creek, Humboldt County, CA.	24
Figure 10. Comparison of 2+ steelhead trout population abundance by month in YR 2011 with the previous seven year average, lower Redwood Creek, Humboldt County, CA.	25
Figure 11. 0+ coho salmon population abundance estimates (error bars are 95% confidence intervals) in YRS 2006 - 2011. Lack of 95% CI for YRS 2009 and 2010 is due to scale of Y axis. Numeric values next to box represent number	

of individuals. Line of best fit is a regression line, with corresponding equation, correlation value (r), and p value, lower Redwood Creek, Humboldt County, CA.	27
Figure 12. Comparison of 0+ coho salmon population abundance by month in YR 2011 with the previous five year average, lower Redwood Creek, Humboldt County, CA.	28
Figure 13. 1+ coho salmon population abundance estimates (error bars are 95% confidence interval) in YRS 2004 – 2011. Lack of 95% CI for YR 2010 is due to scale of Y axis. Numeric values next to box represent number of individuals. Line of best fit is a regression line, with corresponding equation, correlation value (r), and p value, lower Redwood Creek, Humboldt County, CA.	29
Figure 14. Comparison of 1+ coho salmon population abundance by month in YR 2011 with the previous seven year average, lower Redwood Creek, Humboldt County, CA.	30
Figure 15. Cutthroat trout population abundance estimates in YRS 2006 – 2008 and YRS 2010 - 2011 (error bars are 95% confidence intervals). Numeric values next to box represent number of individuals. Line of best fit is a regression line, with corresponding equation, correlation value (r), and p value, lower Redwood Creek, Humboldt County, CA.	32
Figure 16. Comparison of cutthroat trout population abundance by month in YR 2011 with the average of YRS 2006 – 2008 and 2010, lower Redwood Creek, Humboldt County, CA.	32
Figure 17. Average, minimum, and maximum stream temperature (°C) in lower Redwood Creek, Humboldt County, CA., 2011.	44
Figure 18. Average daily stream temperature (°C) in YRS 2004 – 2011, lower Redwood Creek, Humboldt County, CA.	44

LIST OF TABLES

Table 1. Miscellaneous species captured by the smolt trap in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.	13
Table 2. The estimated catch and expansion (population level) of juvenile anadromous salmonids considered to have been missed due to trap not being deployed (n = 5 d) during the emigration period of April 8 through September 9 (as a percentage of total without missed days catch in parentheses), lower Redwood Creek, Humboldt County, CA., 2011.	14
Table 3. Average weekly and seasonal trapping efficiencies for 0+ Chinook salmon, 1+ Chinook salmon, 1+ steelhead trout, 2+ steelhead trout, 0+ coho salmon, 1+ coho salmon, and cutthroat trout in YR 2011, lower Redwood Creek, Humboldt County, CA.	16

Table 4. Date of peak weekly 0+ Chinook salmon population emigration by study year (number of individuals in parentheses), lower Redwood Creek, Humboldt County, CA.	19
Table 5. Production of 0+ Chinook salmon partitioned into fry and fingerling categories each study year and for the previous seven year average (expressed as a percentage in parentheses for YR 2011 and the previous seven year average), lower Redwood Creek, Humboldt County, CA.	20
Table 6. Date of peak weekly 1+ steelhead trout population emigration by study year (number of individuals in parentheses), lower Redwood Creek, Humboldt County, CA.	23
Table 7. Date of peak weekly 2+ steelhead trout population emigration by study year (number of individuals in parentheses), lower Redwood Creek, Humboldt County, CA.	26
Table 8. Date of peak weekly 0+ coho salmon population emigration by study year (number of individuals in parentheses), lower Redwood Creek, Humboldt County, CA.	28
Table 9. Date of peak weekly 1+ coho salmon population emigration by study year (number of individuals in parentheses), lower Redwood Creek, Humboldt County, CA.	31
Table 10. Date of peak weekly cutthroat trout population emigration by study year (number of individuals in parentheses), lower Redwood Creek, Humboldt County, CA.	33
Table 11. Comparison of 0+ steelhead trout, 1+ steelhead trout, and 2+ steelhead trout percent composition of total juvenile steelhead trout downstream migration in YR 2011 with the previous seven year average, lower Redwood Creek, Humboldt County, CA.	34
Table 12. 0+ Chinook salmon average and median fork length (mm) and weight (g) in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.	35
Table 13. 1+ Chinook salmon average and median fork length (mm) and weight (g) in YRS 2005, 2008 - 2011, lower Redwood Creek, Humboldt County, CA.	35
Table 14. 0+ steelhead trout average and median fork length in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.	36
Table 15. 1+ steelhead trout average and median fork length (mm) and weight (g) in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.	37
Table 16. 2+ steelhead trout average and median fork length (mm) and weight (g) in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.	37
Table 17. 0+ coho salmon average and median fork length (mm) and weight (g) in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.	38
Table 18. 1+ coho salmon average and median fork length (mm) and weight (g) in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.	39
Table 19. Cutthroat trout average and median fork length (mm) and weight (g) in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.	39
Table 20. Developmental stages of captured 1+ and 2+ steelhead trout in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.	40
Table 21. Trapping mortality for juvenile salmonids captured in YR 2011, lower Redwood Creek, Humboldt County, CA.	41

Table 22. Comparison of trapping mortality of juvenile salmonids in eight consecutive study years, lower Redwood Creek, Humboldt County, CA.	41
Table 23. Average, minimum, and maximum stream temperatures (°C) (standard error of mean in parentheses) at the trap site during the trapping periods in YRS 2004 – 2011, lower Redwood Creek, Humboldt County, CA.	42
Table 24. Average monthly stream temperature (°C) (°F in parentheses) at the trapping site in study years 2004 - 2011, lower Redwood Creek, Humboldt County, CA.	43
Table 25. Maximum weekly average temperature (MWAT) and maximum weekly maximum temperature (MWMT) for stream temperatures °C (°F in parentheses) at the trap site in lower Redwood Creek, Humboldt County, CA., study years 2004 – 2011.	43

LIST OF APPENDICES

Appendix 1. Reasons for collecting genetic samples from Chinook salmon, steelhead trout smolts, and coho salmon fry, parr, and smolts.	74
Appendix 2. Graphical representation of daily stream gage height (ft.) at trap site and (preliminary) average daily streamflow (cfs) measured at Orick gaging station (USGS 2011), lower Redwood Creek, Humboldt County, CA.	75

2011 CDFG FRGP REPORT

LOWER REDWOOD CREEK JUVENILE SALMONID (SMOLT) ABUNDANCE PROJECT 2004 - 2011 Seasons PROJECT 2a7^{1/}

Prepared by

Michael D. Sparkman
Northern Region

ABSTRACT

Juvenile anadromous salmonid trapping was conducted for the eighth consecutive year in YR 2011 in lower Redwood Creek, Humboldt County, California during the spring/summer emigration period (April – September). Trapping in YR 2011 was continued into September to account for delays in migration. The purpose of the study was to describe juvenile salmonid out-migration and estimate smolt population abundances for wild 0+ Chinook salmon, 1+ Chinook salmon, 0+ coho salmon, 1+ coho salmon, 1+ steelhead trout, 2+ steelhead trout, and cutthroat trout using mark/recapture methods. The long term goal is to monitor the status and trends of out-migrating juvenile salmonid smolts in Redwood Creek in relation to watershed conditions and restoration activities in the basin, and to provide data for Viable Salmonid Population Analysis. A rotary screw trap and fyke net/pip trap collectively operated 149 out of 154 days/nights possible, and captured 56,911 0+ Chinook salmon (ocean type), 11 1+ Chinook salmon (stream type), 9,864 0+ steelhead trout, 2,680 1+ steelhead trout, 721 2+ steelhead trout, 52 cutthroat trout, zero 0+ pink salmon, 226 0+ coho salmon, and 24 1+ coho salmon to total 70,489 juvenile salmonids. Average weekly trapping efficiencies were 36% for 0+ Chinook salmon, 38% for 1+ Chinook salmon, 14% for 1+ steelhead trout, 9% for 2+ steelhead trout, 27% for cutthroat trout, 22% for 0+ coho salmon, and 3% for 1+ coho salmon. The 0+ Chinook salmon population abundance in YR 2011 equaled 147,719 individuals (95% CI = 136,199 – 159,238), and for 1+ Chinook salmon equaled 15 individuals (95% CI = 10 – 20). The low abundance of yearling Chinook salmon indicated they are relatively rare in Redwood Creek. Population abundances with 95% confidence intervals in YR 2011 equaled 20,5011 (17,021 – 23,980) for 1+ steelhead trout; 5,587 (4,094 – 7,081) for 2+ steelhead trout; 884 (639 – 1,128) for 0+ coho salmon; 113 (16 – 209) for 1+ coho salmon, and 118 (73 – 164) for cutthroat trout. The abundance of 1+ coho salmon smolts in YR 2011 was alarmingly low. The population abundances of 0+ Chinook salmon, 0+ coho salmon, 1+ coho salmon, and cutthroat trout each showed a (preliminary) non-significant trend over study years ($p > 0.10$), and for 1+ and 2+ steelhead trout significant, negative trends were detected ($p < 0.10$). The trend in abundance for 0+ Chinook salmon was significantly negative over time when flood type flows in the upper basin were added to the linear model ($p < 0.10$), which indicated that Chinook salmon populations passing through the lower basin were influenced by the Chinook salmon population abundance and flood type flows in the upper basin. The abundances of 1+ and 2+ steelhead trout over time were also negatively related to flood type flows in the upper basin. Monthly peaks in population abundances in YR 2011 occurred later than usual for 0+ Chinook salmon, 1+ Chinook salmon, 1+ steelhead trout, 2+ steelhead trout, 0+ coho salmon, and cutthroat trout, and may be attributable to higher than average stream flows during the trapping period. The two most important months for migration in YR 2011 were June/July for 0+ Chinook salmon, 1+ steelhead trout, 2+ steelhead trout, and 0+ coho salmon; May/June for 1+ Chinook salmon, April/May for 1+ coho salmon, and July/August for cutthroat trout. Considerably more 1+ steelhead trout emigrated downstream than 2+ steelhead trout each study year, and may indicate that stream habitat conditions are limiting the abundance of the older age class, or favoring a change in the life history to a younger smolt age.

^{1/} This paper should be referenced as: Sparkman MD. 2012. Lower Redwood Creek juvenile salmonid (smolt) abundance project, study year 2011: a report to the Fisheries Restoration Grants Program (Project No. P0810509). CDFG AFRAMP, study 2a7: 81 p.

INTRODUCTION

This report presents results of the eighth consecutive year of juvenile salmonid downstream migrant trapping in lower Redwood Creek, Orick, California during the spring/summer emigration period of YR 2011. The study was conducted by the California Department of Fish and Game's Anadromous Fisheries Resource Assessment and Monitoring Program (CDFG AFRAMP) in YRS 2004 – 2011, and is a cooperative effort between the department, United States Geological Survey, Humboldt State University, and the landowner where the trap is located. Funding for YR 2004 was provided by the department's Steelhead Report Card Program and AFRAMP, and in YR 2005 funding was provided by the Steelhead Report Card Program, AFRAMP, and the Fisheries Restoration Grants Program (FRGP). In YRS 2006 – 2007 funding was provided by AFRAMP and FRGP, and in YR 2008 provided by CDFG AFRAMP, CDFG Steelhead Trout Report-Restoration Card Program, FRGP, and Save the Redwoods League. In YRS 2009 - 2011 funding was provided by CDFG AFRAMP and FRGP (YR 2011: P0810509, S02067).

The initial impetus for this study was to determine how many wild salmon and steelhead trout smolts were emigrating from the majority of the Redwood Creek basin before entering the Redwood Creek estuary and Pacific Ocean. The 'majority' of the Redwood Creek basin includes all anadromous waters upstream of the first major tributary (Prairie Creek, river mile RM 3.7) to Redwood Creek. Areas downstream of Prairie Creek are generally not used for spawning by adult salmonids; thus, the only smolt production the trap will miss is from the Prairie Creek watershed. In YR 2004, CDFG AFRAMP successfully determined juvenile Chinook salmon and steelhead trout smolt population abundances from the majority of Redwood Creek for the first time in Redwood Creek's anadromous salmonid monitoring history. Additionally, AFRAMP and the Redwood Creek Landowners Association (RCLA) have successfully determined smolt population abundances for juvenile Chinook salmon and steelhead trout emigrating from upper Redwood Creek for the past twelve consecutive years (Sparkman, In progress). Prior to our studies on juvenile salmonid downstream migration and smolt abundance in Redwood Creek, scientific studies which quantified anadromous salmonids within the Redwood Creek watershed were primarily limited to the estuary (juveniles) and Prairie Creek (adults and juveniles).

Adult salmon and steelhead trout populations are difficult to monitor in Redwood Creek because the adult fish migrate upstream during fall or late fall, winter, and early spring. Thus, when the adults are present, the streamflow is often high and unpredictable, which limits the reliability and usefulness of any adult weir. Additionally, the streamflow during this time period often carries large amounts of suspended sediments, which render visual observations of adult fish and redds (e.g. spawning surveys) unreliable and unlikely for long term monitoring, particularly in average or above average water years. However, efforts are currently underway to count adult fish migrating upstream in lower Redwood Creek with a DIDSON unit (USGS California Cooperative Fish and Wildlife Research Unit at Humboldt State University), and to count redds in randomly selected areas within the Redwood Creek watershed (CDFG AFRAMP). Scientific studies which

focus on adult salmonids in tributaries to Redwood Creek are less affected by these processes (high, muddy stream flow), however, the tributaries are less likely to adequately represent or account for the majority of the salmonid populations in Redwood Creek because the majority of adult salmon and steelhead trout spawn in the mainstem. An exception is the Prairie Creek watershed which probably accounts for a considerable amount of the coho salmon and cutthroat trout production in Redwood Creek. Tributaries to Redwood Creek are often steep, with limited anadromy (RNP 1997, Brown 1988). Additionally, some of the tributaries can dry up prior to late summer, which cause the juvenile fish to migrate into the mainstem of Redwood Creek.

Determining and tracking smolt numbers over time is an acceptable, useful, and quantifiable measure of salmonid populations which many agencies (both state and federal), universities, consultants, tribal entities, and timber companies perform each year. Juvenile salmonid out-migration can be used to assess: 1) the number of parents that produced the cohort (Roper and Scarnecchia 1999, Ward 2000, Sharma and Hilborn 2001, Ward et al. 2002, Bill Chesney pers. comm. 2006), 2) redd gravel conditions (Cederholm et al. 1981, Holtby and Healey 1986, Hartman and Scrivener 1990), 3) in-stream habitat quality and watershed health (Tripp and Poulan 1986, Hartman and Scrivener 1990, Hicks et al. 1991, Bradford et al. 2000, Sharma and Hilborn 2001, Ward et al. 2002), 4) restoration activities (Everest et al. 1987 *in* Hicks et al. 1991, Slaney et al. 1986, Tripp 1986, McCubbing and Ward 1997, Solazzi et al. 2000, Cleary 2001, Ward et al. 2002, McCubbing 2002, Ward et al. 2003, Roni et al. 2006), 5) over-winter survival (Scrivener and Brown 1993 *in* McCubbing and Ward 1997, Quinn and Peterson 1996, Solazzi et al. 2000, McCubbing 2002, Ward et al. 2002, Giannico and Hinch 2003, Ebersole et al. 2009), and 6) future recruitment to adult populations (Holtby and Healey 1986, Nickelson 1986, Ward and Slaney 1988, Ward et al. 1989, Unwin 1997, Ward 2000).

This paper will present results of trapping in YR 2011 with various comparisons to the average of the previous seven years (YRS 2004 – 2010), and YR 2010.

Site Description

Redwood Creek lies within the Northern Coast Range of California, and flows 67 miles through Humboldt County before reaching the Pacific Ocean (Figure 1). Headwaters originate at an elevation of about 5,000 ft and converge to form the main channel at about 3,100 feet. Redwood Creek flows north to northwest to the Pacific Ocean, and bisects the town of Orick in Northern California. The basin of Redwood Creek is 179,151 acres, about 49.7 miles long, and 6.2 miles wide (Cashman et. al 1995). The study area upstream of the trap site (Rm 4) encompasses approximately 151,922 acres of the Redwood Creek basin, with about 93 stream miles (150 km) of accessible salmon and steelhead habitat (Cannata et al. 2006).




Figure 1. Redwood Creek basin with rotary screw trap location (RM 4), Humboldt County, CA. (scale is slightly inaccurate due to reproduction process, Charlotte Peters pers. com. 2001).

Geology

The Redwood Creek watershed is situated in a tectonically active and geologically complex area, and is considered to have some of the highest uplift and seismic activity rates in North America (CDFG NCWAP 2004).

The geology of the Redwood Creek basin has been well-studied and mapped (Cashman et. al 1995).

“Redwood Creek drainage basin is underlain by metamorphic and sedimentary rocks of the Franciscan assemblage of Late Jurassic and Early Cretaceous age and by shallow marine and alluvial sedimentary deposits of late Tertiary and Quaternary age. These units are cut by a series of shallowly east-dipping to vertical north to northwest trending faults. The composition and distribution of bedrock units and the distribution of major faults have played a major part in the geomorphic development of the basin. Slope profiles, slope gradients, and drainage patterns within the basin reflect the properties of the underlying bedrock. The main channel of Redwood Creek generally follows the trace of the Grogan fault, and other linear topographic features are developed along major faults. The steep terrain and the lack of shear strength of bedrock units are major contributing factors to the high erosion rates in the basin” (Cashman et al. 1995).

Climate

The climate of the Redwood Creek basin varies dependent upon location within the watershed and season. Coastal areas have a moderate climate due to proximity to the ocean, and differ from inland areas (i.e. upper Redwood Creek) which experience higher and lower temperatures. Summers are typically cool and moist on the coast, and hot and dry inland. Snowfall is common during winter months in the upper basin and relatively rare in the lower basin.

Stream Discharge

A USGS gauging station (#11482500) is located about 850 m downstream of the trap site in lower Redwood Creek. The gauging station is downstream of the confluence of Prairie Creek with Redwood Creek, thus the station is influenced by Prairie Creek streamflow. Streamflow records for the Orick gage cover the periods of 1911 – 1913, 1953 – 2011, and total 60 years (USGS 2011). High streamflows usually occur from November through May, and typically peak in January. However, the months of December, February, March, and April can experience peaks in high flows as well. Using all years' data (historic), mean monthly discharge was 1,008 cfs (28.5 m³/sec), and ranged from 35 – 2,503 cfs (0.99 – 70.9 m³/sec) (USGS 2011). Average monthly discharge in WY 2011 (preliminary data) equaled 1,119 cfs (31.7 m³/sec), ranged from 20 – 3,937 cfs (0.6 – 111.5 m³/sec), and peaked in March. Average streamflow in WY 2011 (preliminary) was about 1.1 times greater than the historic average, and the second highest of the eight current study years.

The 60 year average monthly flow during the majority of the (normal) trapping season (April – July) equaled 557 cfs (15.8 m³/sec), and ranged from 86 – 1,244 cfs (2.4 – 35.2 m³/sec) (USGS 2011). Average monthly discharge from April – July in YR 2011 (preliminary data) equaled 757 cfs (21.4 m³/sec), ranged from 106 – 1,982 cfs (3.0 – 56.1 m³/sec), and was 1.36 times greater than the historic average for this time frame (USGS 2011). Due to the increase in discharge in YR 2011, trapping was extended until September 9th in order to fully account for the additional migration which occurred in August and early September.

Overstory

The overstory of Redwood Creek is predominately second and third growth Redwood (*Sequoia sempervirens*) and Douglas Fir (*Pseudotsuga menziesii*), mixed with Big Leaf Maple (*Acer macrophyllum*), California Bay Laurel (*Umbellularia californica*), Incense Cedar (*Calocedrus decurrens*), Cottonwood (*Populus spp.*), Manzanita (*Arctostaphylos spp.*), Oak (*Quercus spp.*), Tan Oak (*Lithocarpus densiflorus*), Pacific Madrone (*Arbutus menziesii*), and Red Alder (*Alnus rubra*). The lower portion of Redwood Creek (ie within Redwood National Park boundaries) contains old growth Redwood, mixed with second growth redwood and other tree species.

Understory

Common understory plants include: dogwood (*Cornus nuttallii*), willow (*Salix lucida*), California hazelnut (*Corylus rostrata*), lupine (*Lupinus spp.*), blackberry (*Rubus spp.*), plantain (*Plantago coronopus*), poison oak (*Toxicodendro diversilobum*), wood rose (*Rosa gymnocarpa*), false Solomon's seal (*Smilacina amplexicaulis*), spreading dog bane (*Apocynum spp.*), wedgeleaf ceanothus (*Ceanothus spp.*), bracken fern (*Pteridium aquilinum*), blackcap raspberry (*Rubus spp.*), and elderberry (*Sambucus spp.*), among other species.

Redwood Creek History (Brief)

Redwood Creek watershed has experienced extensive logging of Redwood and other commercial tree species. By 1978, 81% of the original forest was logged, totaling 66% of the basin area (Kelsey et al. 1995). Most, if not all, remaining old growth Redwood is contained within Redwood National Park, which is about 200 m upstream of the trap site. In conjunction with clear-cut logging, log removal via tractors, associated road building, geology types and geomorphic processes (eg debris slides and earthflows), and flood events in 1955 and 1964, large amounts of sediments were delivered into the stream channel (Madej and Ozaki 1996) with a resultant loss of stream habitat complexity (filling in of pools and flattening out of the stream channel, Marlin Stover pers. comm. 2000). Additional high flows occurred in 1972, 1975, and 1995 as well, and have helped influence the current channel morphology of Redwood Creek. The downstream migrant trap in lower Redwood Creek is located in an area of gravel aggradation, and gravel extraction does occur in this area. Redwood Creek has been listed as sediment and

temperature-impaired under section 303(d) of the Clean Water Act (CWA 2002; SWRCB 2003; USEPA 2003).

Federal ESA Species Status

Chinook (King) salmon (*Oncorhynchus tshawytscha*), coho (Silver) salmon (*O. kisutch*), steelhead trout (*O. mykiss*), and cutthroat trout (*O. clarki clarki*) are known to inhabit Redwood Creek. This study and the study in upper Redwood Creek also show that pink salmon (*O. gorbuscha*) are present in Redwood Creek. Chinook salmon (KS) of Redwood Creek belong to the California Coastal Chinook Salmon Evolutionarily Significant Unit (ESU), and are listed as “threatened” under the Federal Endangered Species Act (Federal Register 1999a). The definition of threatened as used by National Oceanic and Atmospheric Administration (NOAA) and the National Marine Fisheries Service (NMFS) is “likely to become endangered in the foreseeable future throughout all or a significant portion of their range” (NOAA 1999). Coho salmon (CO) belong to the Southern Oregon/Northern California Coasts ESU and were classified as “threatened” (Federal Register 1997) prior to the Chinook salmon listing. Steelhead trout (SH) fall within the Northern California Steelhead ESU, and are also listed as a “threatened” species (Federal Register 2000). Coastal cutthroat trout (CT) of Redwood Creek fall within the Southern Oregon/California Coasts Coastal Cutthroat Trout ESU, and were determined “not warranted” for ESA listing (Federal Register 1999b). Despite ESU listings of Redwood Creek anadromous salmonid populations, relatively little data exists concerning abundance and population sizes, particularly for juvenile (and adult) life history stages. Historically, the most prolific species was most likely the fall/early winter-run Chinook salmon.

Purpose

The purpose of this project is to describe juvenile salmonid downstream migration from the majority of the Redwood Creek basin, and to determine emigrant population abundances for wild 0+ (young-of-year) Chinook salmon (ocean type), 1+ Chinook salmon (stream type) (between 1 and 2 years old), 1+ steelhead trout, 2+ (2 years old and greater) steelhead trout, cutthroat trout (age 1 and older), 0+ coho salmon (fry, parr), and 1+ coho salmon smolts. The primary long term goal is to monitor the status and trends of out-migrating juvenile salmonid smolts in Redwood Creek in relation to watershed condition and restoration activities in the basin; and to provide data needed for Viable Salmonid Population (VSP) analysis. Specific study objectives were as follows:

- 1) Determine the specie composition and temporal pattern of downstream migrating juvenile salmonids.
- 2) Determine population estimates for downstream migrating 0+ Chinook salmon, 1+ Chinook salmon, 1+ steelhead trout, 2+ steelhead trout, cutthroat trout, 0+ coho salmon, and 1+ coho salmon.
- 3) Record fork length (mm) and weight (g) of captured fish.

- 4) Collect genetic samples from 0+ Chinook salmon, 1+ Chinook salmon, 0+ steelhead trout, 1+ steelhead trout, 2+ steelhead trout and juvenile coho salmon (if present) for future analyses and comparisons (Appendix 1).
- 5) Collect and handle fish in a manner that minimizes mortality.
- 6) Statistically analyze data for significance and trends.
- 7) Compare data between study years.
- 8) Link data collected from the lower trap, upper trap, and estuary (Redwood National Park) to provide a more complete study on the life history and abundance of emigrating juvenile salmonids (smolts) in Redwood Creek.

METHODS AND MATERIALS

Trap Operations

The methods and materials used in this study in YR 2011 were the same as previous study years (Sparkman 2011a). A modified E.G. Solutions (5 foot diameter cone) rotary screw trap was deployed in lower Redwood Creek (Rm 4) on April 8th, 2011 at the same general location (\pm 100 m) as previous study years (YRS 2004 - 2010).

We operated the rotary screw trap continually (24 hrs/day, 7 days a week) from April 8th through September 2nd, with exception to five days of missed trapping. Five days were missed on April 16 – April 20 when stream flow and potential debris loading in the trap's livebox were too high to safely trap. Beyond September 2nd, stream flow was too low to operate the rotary screw trap, and on September 2nd we installed a fyke net/pipe trap. Weir panels were used to force all downstream migrating fish into the fyke net/pipe trap. The trapping season in YR 2011 was extended compared to previous study years because smolt migrations in August were much higher than expected. Trapping was discontinued on September 9th when the catch distribution for each species at age reached zero, or when relatively few individuals were caught in consecutive days.

During periods of lesser streamflows, weir panels were used with the rotary screw trap to: 1) keep the trap's cone revolutions relatively high, and 2) maintain good trapping efficiencies by directing fish into the cone area. The weir panels were held in place using bailing wire and 6 - 8 ft long fence posts, and were first installed in May. Additional weir panels were later added to increase the overall length, and by July 13th the panels reached the pontoons of the smolt trap.

The YR 2011 trapping season can be characterized as: working in and out of high stream flows in April, making frequent adjustments to the trap configuration, maintaining the trap's position in the thalweg, and extensively using weir panels. The largest flow event occurred on April 16th, 2011 when the stream rose from 745 cfs (on April 15th) to 5,020 cfs (April 16th).

Trapping in YR 2011 was less difficult than trapping in YR 2010. Adjustments to trap placement and the use of weir panels in YR 2011 helped increase trapping efficiencies and cone revolutions.

Biometric Data Collection

Fishery technicians frequently removed debris (e.g. alder cones, leaves, sticks, detritus, etc.) from within the livebox at night to reduce trap mortalities the following morning. The trap's livebox was emptied at 09:00 every morning by 2 - 4 technicians. Debris was once again inspected and carefully removed so that the smaller fish would not be released into the stream with the debris.

Young of year fish were removed first and processed before 1+ and 2+ fish to decrease predation or injury to the smaller fish. Captured fish (0+ fish first, then 1+ and older) were placed into 5 gal. buckets and carried to the processing station. At the station, fish were placed into a 23.5 gal. ice chest modified to safely hold juvenile fish. The ice chest was adapted to continually receive fresh water from the stream using a 3,700 gph submersible bilge pump. The bilge pump connected to a flexible line (ID 4 cm or 1.6 in.) that connected to a manifold with four ports. "Y" type hose adapters were connected to each port. Garden hoses connected to the hose adapters, with one line feeding the ice chest, and four lines feeding recovery buckets for processed fish. Additional garden hoses were connected to the hose adaptors to quickly fill buckets if needed, and to relieve any excess back pressure. Plumbing inside the ice chest consisted of two PVC pipes: one that served to dissipate the stream water into the ice chest, and the other to adjust water height in the ice chest and to drain excess water. The system worked very well, did not require additional battery operated aerators, and decreased total fish processing time.

Random samples of each species at age (eg 0+ KS, 0+ SH, etc.) were netted from the ice chest for examination, enumeration, and biometric data collection. Each individual fish was counted by species at age, and observed for trap efficiency trial marks. Marked fish from the trap in upper Redwood Creek (Sparkman 2011b) were tallied separately from the marked fish used to determine trapping efficiencies for the lower trap.

Fork Lengths/Weights

Fish were anesthetized with MS-222 prior to data collection in 2 gal. dishpans. Biometric data collection included 30 measurements of fork length (mm) and wet weight (g) for random samples of 0+ Chinook salmon (0+ KS), 1+ Chinook salmon (1+ KS, if present), 1+ and greater cutthroat trout (CT), 1+ steelhead trout (1+ SH), 2+ and greater steelhead trout (2+ SH), 0+ coho salmon (0+ CO), 1+ coho salmon (1+ CO), and 0+ pink salmon (if present). Only fork lengths were taken from 0+ steelhead trout (0+ SH). A 160 and 350 mm measuring board (± 1 mm), and an Ohaus Scout II digital scale (± 0.1 g) were used in the study. Fork lengths were taken every day of trap operation, and fork length frequencies of 0+ and older steelhead trout, coho salmon, and Chinook salmon were used

to determine age-length relationships at various times throughout the trapping period. Scales were occasionally read to verify age class cutoffs. 0+ Chinook salmon and 1+ steelhead trout weights were taken 2 - 7 times per week; and 0+ and 1+ coho salmon, and 2+ steelhead trout weights were taken nearly every day of trap operation and collection due to expected, low sample sizes. Individuals were weighed in a tared plastic pan (containing water) on the electronic scale. The scale was placed in a large plastic bin when weighing fish to prevent any influences from wind, and was calibrated every day prior to data collection. After biometric data was collected, fish were placed into 5 gal. recovery buckets which received continuously pumped fresh stream water. Young of year fish were kept in separate recovery buckets from age 1+ and older fish to decrease predation or injury. When fully recovered from anesthesia, 0+ juvenile fish were transported 150 m downstream of the trap site and released in the margin of the stream; and aged 1 and older fish were transported 200 m downstream of the trap site and released near the middle of the stream when possible.

Population Estimates

The number of fish captured by the trap represented only a portion of the total fish moving downstream in that time period. Total salmonid out-migration estimates (by age and species) were determined on a weekly basis for 0+ Chinook salmon, 1+ Chinook salmon, 1+ steelhead trout, 2+ steelhead trout, cutthroat trout, 0+ coho salmon, and 1+ coho salmon using stratified and non-stratified mark-recapture methods described by Carlson et al. (1998). Population estimation methods in YR 2011 were identical to previous study years (Sparkman 2011a). Annual variation in both population abundances and catches over the current eight year period were characterized by the standard deviation and standard error of the mean for each species at age.

Physical Data Collection

A staff gage with increments in hundredths of a foot was used to measure the relative stream surface elevation (hydrograph) at the trap site each day from April 9th – September 9th, 2011. A graphical representation of the data, along with average daily stream discharge data (preliminary) from the Orick gaging station (USGS 2011), is given in Appendix 2. Stream temperatures were recorded with an Optic StowAway® Temp data logger (Onset Computer Corporation, 470 MacArthur Blvd. Bourne, MA 02532) placed behind the rotary screw trap. The probe was placed into a PVC cylinder with holes to ensure adequate ventilation and to prevent influences from direct sunlight. The probe recorded stream temperatures (°C) every 30 minutes and recorded 7,393 measurements over the course of the study. The shallowest stream depth during which measurements were taken (in September) was about 1.5 feet.

Statistical Analyses

The statistical analyses conducted in YR 2011 were the same as in previous study years (Sparkman 2011a). Numbers Cruncher Statistical System software (NCSS 97) (Hintze 1998) was used for linear correlation, regression/ANOVA output, and descriptive statistics. Linear regression was used to estimate the catch for each species at age for days when the trap was not fishing by using data before and after the missed day(s) catch. The estimated catch (except for 0+ steelhead) was then added to the known catch in a given stratum and applied to the population model for that stratum (Roper and Scarnecchia 1999). Linear correlation slope and equation line were used to determine if total catches and population abundances of a given species at age were increasing or decreasing over the eight years of study. The tests are considered preliminary, and more data will be required to detect the true trends in population abundances over years. With respect to 0+ Chinook salmon, peaks in stream flows were great enough to potentially mobilize redd gravels each study year. Flood type flows capable of gravel scour (and deposition) in mid to lower Redwood Creek are generally thought to occur near 11,000 cfs (Randy Klein, Greg Bundros, Vicki Ozaki, Mary Ann Madej, pers. comm. 2003). Peak winter flows in upper Redwood Creek, coded as 1 or 0, were included in additional correlation tests with study year on population size for 0+ Chinook salmon, and 1+ and 2+ steelhead trout passing through the lower basin. High bedload mobilizing flows were coded as 1 (for population estimates in YRS 2005 and 2006) and non-bedload mobilizing flows as 0 (for population estimates in YRS 2004, 2007, 2008, 2009, 2010, and 2011) (Zar 1999). The test for 0+ Chinook salmon would indicate if the relationship of peak winter flows during egg incubation in spawning redds in the upper basin decreased survival, and hence impact the numbers migrating downstream, and tests for 1+ and 2+ steelhead trout would indicate if high winter flows were affecting population abundances of steelhead smolts from upper Redwood Creek with respect to over-winter survival. Flows considered great enough to mobilize the bedload in upper Redwood Creek (> 6,000) were identified by Redwood National Park hydrologists and Geologists), and based upon smolt data collected in the upper basin, appear to represent a threshold for survival.

Descriptive statistics were used to characterize the average FL (mm) and Wt (g) of each species at age on a study year basis. Linear regression was used to test the relationship of average daily discharge on average daily stream temperature, and correlation was used to test whether the average daily stream temperature increased over time (d) during the study period. If data violated tests of statistical assumptions (n = 4 tests for ANOVA, n = 3 tests for regression/correlation; NCSS 97), data was transformed with Log (x +1) to approximate normality (Zar 1999). The term 'transformed' in this paper refers to the log (x +1) transformation. Power is defined as the probability of correctly rejecting the null hypothesis when it is false; and can also be thought of as the probability of detecting differences that truly exist (Zar 1999). The level of significance (alpha) was set at 0.10 for trend analysis purposes.

RESULTS

The rotary screw trap operated from 4/08/11 – 9/02/11 and trapped 142 days/nights out of a possible 147, and the fyke net/pipe trap operated from 9/02/11 – 9/09/11 and trapped 7 days/nights out of a possible 7. The five missed days of trapping occurred from 4/16/11 – 4/20/11. The trapping rate in YR 2011 was 97%, compared to 96% for the previous seven year average (ranged from 91 – 99%).

Species Captured

Juvenile Salmonids

Species captured in YR 2011 included: juvenile Chinook salmon (*Oncorhynchus tshawytscha*), juvenile coho salmon (*O. kisutch*), juvenile steelhead trout (*O. mykiss*), and coastal cutthroat trout (*O. clarki clarki*). A total of 70,489 juvenile salmonids were captured in YR 2011 (Figure 2).




Figure 2. Total juvenile salmonid trap catches (n = 70,489) from April 9th through September 9th, 2011, lower Redwood Creek, Humboldt County, CA. Numeric values above columns represent actual catches. 0+ KS = young-of-year Chinook salmon, 1+ KS = age 1 Chinook salmon, 0+ SH = young-of-year steelhead trout, 1+ SH = age 1 and older steelhead trout, 2+ SH = age 2 and older steelhead trout, 0+ CO = young of year coho salmon, 1+ CO = age 1 and older coho salmon, CT = cutthroat trout, 0+ Pink = young-of-year pink salmon.

The average total catch by study year (n = 8 years) equaled 70,348 (SD = 33,192; SEM = 11,735). The eight year average catch equaled 45,512 (SD = 22,304; SEM = 7,886) for 0+ Chinook salmon, 13 (SD = 20; SEM = 7) for 1+ Chinook salmon, 18,698 (SD = 16,902; SEM = 5,976) for 0+ steelhead trout, 5,044 (SD = 2,225; SEM = 787) for 1+ steelhead trout, 806 (SD = 368; SEM = 130) for 2+ steelhead trout, 164 (SD = 137; SEM = 48) for 0+ coho salmon, 75 (SD = 74; SEM = 26) for 1+ coho salmon, 36 (SD = 24; SEM = 9) for cutthroat trout, and 0.3 (SD = 0.8; SEM = 0.3) for 0+ pink salmon.

Miscellaneous Species

The trap caught numerous species besides juvenile anadromous salmonids in YR 2011, including: prickly sculpin (*Cottus asper*), coast range sculpin (*Cottus aleuticus*), sucker (*Catostomidae* family), three-spined stickleback (*Gasterosteus aculeatus*), juvenile (ammocoete) lamprey and adult Pacific Lamprey (*Lampetra tridentatus*), and western Brook Lamprey (*Lampetra richarsoni*), among other species (Table 1). Adult and juvenile captures occurred for Prickly Sculpin, Coast Range Sculpin, Sucker, 3-Spined Stickleback, and Pacific Lamprey. Many gravid sculpins (both species) were also captured. For the second time in eight consecutive years, a bullhead (catfish; *ameiurus* spp.) was captured in YR 2011.

Table 1. Miscellaneous species captured by the smolt trap in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.

Species Captured	YR 2004	YR 2005	YR 2006	YR 2007	YR 2008	YR 2009	YR 2010	YR 2011
Prickly Sculpin	68	140	209	109	234	193	187	491
Coast Range Sculpin	502	212	599	478	3,798	1,576	1,482	969
Sucker	156	89	194	575	121	21	121	51
3-Spined Stickleback	7,225	215	1,119	4,550	1,434	2,176	1,933	808
Bullhead	0	0	0	0	0	0	1	1
Adult Pac. Lamprey	13	3	4	19	16	5	6	4
Juvenile Lamprey*	154	84	50	144	67	50	131	253
Brook Lamprey	0	1	1	0	2	1	1	1
Pac. Giant Salamander	4	8	15	12	8	2	4	16
Painted Salamander	0	0	0	0	0	0	0	0
Rough Skinned Newt	2	3	0	8	16	5	6	4
Red-Legged Frog	0	2	0	4	8	0	11	3
Yellow-Legged Frog	0	0	0	0	0	5	13	5
Tailed Frog**	0	1	0	2	0	0	5	6
Western Toad							161	79
Crawfish								1
Bull Frog								1

* Ammocoete stage. ** Includes adult and tadpole stage.

Days Missed Trapping

Five days were not trapped during the course of study in YR 2011 due to high flow events and potentially, high debris loading in the trap's livebox. The five days of missed trapping would not have influenced the total catch or population estimate of any species at age to any large degree (Table 2).

Table 2. The estimated catch and expansion (population level) of juvenile anadromous salmonids considered to have been missed due to trap not being deployed (n = 5 d) during the emigration period of April 8 through September 9 (as a percentage of total without missed days catch in parentheses), lower Redwood Creek, Humboldt County, CA., 2011.

Age/spp*	Catch	Population Level
0+ KS	205 (0.36%)	3,827 (2.67%)
1+ KS	0 (0.00%)	0 (0.00%)
0+ SH	9 (0.09%)	-
1+ SH	58 (2.16%)	145 (0.75%)
2+ SH	30 (4.34%)	75 (1.34%)
0+ CO	0 (0.00%)	0 (0.00%)
1+ CO	0 (0.00%)	0 (0.00%)
CT	0 (0.00%)	0 (0.00%)

* Age/species abbreviations are the same as in Figure 2.

Note: Regression methods were used to estimate the number of fish caught when the trap was not operating. The estimated catches were then added the known catches for a given stratum (week) and used in the population estimate for that stratum (Roper and Scarnecchia 1999).

Trends in Trap Catches

0+ Chinook Salmon

Linear correlation detected a non-significant, positive relationship of trap catches over study years (n = 8, p = 0.38, r = 0.30, power = 0.13). The correlation of 0+ Chinook salmon trap catches and flood type flows in the upper basin (dummy variable) during egg incubation with study years was significantly negative (n = 8, p = 0.02, Adj. r = 0.83, negative slope for both x variables, power = 0.48). The correlation of trap catches and flood type flows in the upper basin was also significantly negative (n = 8, p = 0.004, R² = 0.77, negative slope, power = 0.96).

1+ Chinook Salmon

Linear correlation detected a non-significant relationship of 1+ Chinook salmon trap catches (transformed) over study years ($n = 8$, $p = 0.21$, $r = 0.50$, positive slope, power = 0.22). The addition of flood type flows in the model did not change the test conclusion ($n = 8$, $p = 0.49$, adj. $r = 0.00$, positive slope for year, negative slope for flood type flows, power = 0.09).

0+ Steelhead Trout

Linear correlation detected a non-significant relationship of 0+ steelhead trout trap catches over study years ($n = 8$, $p = 0.59$, $r = 0.23$, negative slope, power = 0.08).

1+ Steelhead Trout

Linear correlation detected a non-significant relationship of 1+ steelhead trout trap catches over study years ($n = 8$, $p = 0.35$, $r = 0.38$, negative slope, power = 0.14). The correlation of trap catches and flood type flows in the upper basin (dummy variable) with study years was non-significant ($n = 8$, $p = 0.53$, Adj. $r = 0.00$, negative slope for both x variables, power = 0.08). The correlation of trap catches and flood type flows in the upper basin was also non-significant ($n = 8$, $p = 0.89$, $r = 0.05$, negative slope, power = 0.05).

2+ Steelhead Trout

Linear correlation detected a non-significant relationship of 2+ steelhead trout trap catches over study years ($n = 8$, $p = 0.59$, $r = 0.23$, negative slope, power = 0.08). The correlation of trap catches and flood type flows in the upper basin (dummy variable) with study years was non-significant ($n = 8$, $p = 0.78$, Adj. $r = 0.00$, negative slope for both x variables, power = 0.06). The correlation of trap catches and flood type flows in the upper basin was also non-significant ($n = 8$, $p = 0.87$, $r = 0.07$, negative slope, power = 0.05).

0+ Coho Salmon

Linear correlation detected a non-significant relationship of 0+ coho salmon trap catches over study years ($n = 8$, $p = 0.92$, $r = 0.04$, negative slope, power = 0.05).

1+ Coho Salmon

Linear correlation detected a non-significant relationship of 1+ coho salmon trap catches over study years ($n = 8$, $p = 0.90$, $r = 0.06$, negative slope, power = 0.05).

Cutthroat Trout

Linear correlation detected a non-significant relationship of cutthroat trout trap catches over study years ($n = 8$, $p = 0.28$, $r = 0.43$, positive slope, power = 0.17).

Trapping Efficiencies

The average trapping efficiency by week and seasonal trapping efficiency for 0+ Chinook salmon, 1+ Chinook salmon, 1+ steelhead trout, 2+ steelhead trout, 0+ coho salmon, 1+ coho salmon and cutthroat trout fell within the range of 3 to 63% (Table 3).

Table 3. Average weekly and seasonal trapping efficiencies for 0+ Chinook salmon, 1+ Chinook salmon, 1+ steelhead trout, 2+ steelhead trout, 0+ coho salmon, 1+ coho salmon, and cutthroat trout in YR 2011, lower Redwood Creek, Humboldt County, CA.

Study Year	Trapping Efficiency (percentage)	
	Average Weekly	Seasonal
0+ Chinook Salmon	35.5	42.5
1+ Chinook Salmon	37.5	62.5
1+ Steelhead Trout	14.1	13.2
2+ Steelhead Trout	9.0	9.5
0+ Coho Salmon	22.1	22.6
1+ Coho Salmon	2.8	5.3
Cutthroat Trout	26.7	34.0

Population Estimates

0+ Chinook Salmon

The population abundance (or production) of 0+ Chinook salmon emigrating past the trap in lower Redwood Creek in YR 2011 equaled 147,719 individuals with a 95% CI of 136,199 – 159,238 (Figure 3). Population estimate error (or uncertainty) equaled $\pm 7.8\%$. Population abundance in YR 2011 was greater than YRS 2005 – 2007 and YR 2010, and less than emigration in YRS 2004, 2008, and 2009 (Figure 3). Population abundance in YR 2011 was 28% less than the previous seven year average ($N_{\text{avg } 7\text{yr}} = 203,939$). The average population abundance over the current eight year period equaled 196,911 (SD = 148,948; SEM = 52,661). Correlation of time (study year) on yearly population abundance indicated a non-significant, negative relationship ($n = 8$, $p = 0.26$, $r = 0.48$, power = 0.20) (Figure 3). Peaks in streamflows (11,000 cfs) capable of redd scour in lower Redwood Creek occurred each study year. The test of time and bedload mobilizing flows in upper Redwood Cr (O’Kane gaging station) on population abundance showed a significant, negative relationship over years (Regression, $n = 8$, $p = 0.04$, adj. $r = 0.78$, negative slope for both ‘x’ variables, power = 0.38).




Figure 3. 0+ Chinook Salmon population abundance estimates (error bars are 95% confidence interval) in YRS 2004 – 2011. Numeric values next to box represent number of individuals. Line of best fit is a regression line with corresponding equation, correlation value (r), and p value, lower Redwood Creek, Humboldt County, CA.

The pattern in monthly population abundance in YR 2011 showed the temporal delay in migration compared to the previous seven year average (Figure 4). Monthly population emigration peaked in July (N = 75,872 or 51% of total) in YR 2011, and for the previous seven year average peaked in June (N = 102,698 or 50% of total) (Figure 4). In YR 2010, monthly emigration also peaked in July (N = 85,499 or 64%). The two most important months for 0+ Chinook salmon population emigration were June and July (74% of total) in YR 2011, and June and July (71% of total) for the previous seven year average (Figure 4). In YR 2010 July and August were the two most important months, and accounted for 77% of the total population abundance.




Figure 4. Comparison of 0+ Chinook salmon population abundance by month in YR 2011 with the previous seven year average, lower Redwood Creek, Humboldt County, CA.

The peak in weekly population emigration in YR 2011 occurred 6/25 – 7/01, and was earlier than the peak in YR 2010 (Table 4). The peak in YR 2011 was one week later than for the peak for the previous seven year average. The largest weekly peak in abundance occurred in YR 2004 (N = 110,980) and the smallest peak occurred in YR 2011 (N = 27,057) (Table 4). The average FL (mm) for 0+ Chinook salmon migrants during the two modes in migration in YR 2011 equaled 40 mm for 4/23 – 4/29, and 67 mm for 6/25 – 7/01 (Figure 5).

Table 4. Date of peak weekly 0+ Chinook salmon population emigration by study year (number of individuals in parentheses), lower Redwood Creek, Humboldt County, CA.

Study Year	Date of peak in weekly emigration (number in parentheses)
2004	6/18 – 6/24 (110,980)
2005	7/16 – 7/22 (29,766)
2006	6/11 – 6/17 (27,889)
2007	6/18 – 6/24 (38,315)
2008	6/25 – 7/01 (37,976)
2009	6/18 – 6/24 (33,430)
2010	7/16 – 7/22 (34,813)
2011	6/25 – 7/01 (27,057)




Figure 5. 0+ Chinook salmon population abundance by week in YR 2011, lower Redwood Creek, Humboldt County, CA.

0+ Chinook salmon downstream migrants consisted of fry (FL < 45 mm) and fingerlings (FL > 44 mm), and the number and percentage of 0+ Chinook salmon migrants grouped into fry or fingerling categories varied among study years (Table 5). In YR 2011, fry comprised 18.8% and fingerlings comprised 81.2% of the total Chinook salmon population abundance (Table 5).

Table 5. Production of 0+ Chinook salmon partitioned into fry and fingerling categories each study year and for the previous seven year average (expressed as a percentage in parentheses for YR 2011 and the previous seven year average), lower Redwood Creek, Humboldt County, CA.

Study Year	0+ Chinook Salmon production as:	
	Fry (FL < 45mm)	Fingerling (FL > 44 mm)
2004	82,584	472,306
2005	2,052	129,113
2006	71	85,078
2007	3,772	137,287
2008	2,589	171,169
2009	9,839	198,980
2010	11,526	121,207
Avg.	16,062 (7.9)	187,877 (92.1)
YR 2011	27,809 (18.8)	119,910 (81.1)

1+ Chinook Salmon

The population abundance (or production) of 1+ Chinook salmon emigrating past the trap in lower Redwood Creek in YR 2011 equaled 15 individuals with a 95% CI of 10 – 20 (Figure 6). Population estimate error (or uncertainty) equaled 33%. June accounted for the majority of population abundance in YR 2011 (59% of total). May accounted for the majority of abundance in YR 2009 (68%) and in YR 2010 (53% of total). The two most important months were May/June in YRS 2011 and 2010, and April/May in YR 2009.




Figure 6. 1+ Chinook Salmon population abundance estimates (error bars are 95% confidence interval) in YRS 2009 – 2011. Lack of 95% CI for YR 2011 is due to scale of ‘Y’ axis. Numeric values next to box represent number of individuals, lower Redwood Creek, Humboldt County, CA.

1+ Steelhead trout

The population estimate of 1+ steelhead trout emigrating past the trap site in lower Redwood Creek in YR 2011 equaled 20,501 individuals with a 95% CI of 17,021 – 23,980 (Figure 7). Population estimate error (or uncertainty) equaled $\pm 17.0\%$. Population abundance in YR 2011 was the lowest of record, and 53% less than the previous seven year average ($N_{\text{avg } 7 \text{ yr}} = 43,230$). The average in population abundance over the current, eight year period equaled 40,389 (SD = 16,973; SEM = 6,001).

Correlation of time (study year) on yearly population abundances indicated a significant, negative relationship ($n = 8$, $p = 0.03$, $r = 0.75$, power = 0.63) (Figure 7). On average, there were 5,172 less individuals each study year (Figure 7). The test of time and bedload mobilizing flows in upper Redwood Cr (O’Kane gaging station) on population abundance also showed a significant, negative relationship over years (Regression, $p = 0.02$, adj. $r = 0.85$, negative slope for both ‘x’ variables, power = 0.54).




Figure 7. 1+ steelhead trout population abundance estimates (error bars are 95% confidence interval) in YRS 2004 – 2011. Numeric values next to box represent number of individuals. Line of best fit is a regression line, with corresponding equation, correlation value (r), and p value, lower Redwood Creek, Humboldt County, CA.

Monthly population emigration peaked in July in YR 2011 (N = 7,630 or 37% of total), compared to June (N = 15,030 or 35% of total) for the previous seven year average (Figure 8). In YR 2010, monthly emigration peaked in May (N = 7,838 or 29% of total). The two most important months for 1+ steelhead trout population emigration were June and July in YR 2011 (57% of total), compared to May and June for the previous seven year average (69% of total). In YR 2010 May and August were the two most important months, and accounted for 46% of the total population abundance.

The peak in weekly population abundance in YR 2011 occurred 6/25 – 7/01, and occurred later than peaks in previous study years (Table 6). For the eight study years, two peaks occurred in late April/early May, two peaks occurred in May, one in late May/early June, two in June, and one in late June/early July (Table 6).




Figure 8. Comparison of 1+ steelhead trout population abundance by month in YR 2011 with the previous seven year average, lower Redwood Creek, Humboldt County, CA.

Table 6. Date of peak weekly 1+ steelhead trout population emigration by study year (number of individuals in parentheses), lower Redwood Creek, Humboldt County, CA.

Study Year	Date of peak in weekly out-migration (number in parentheses)
2004	5/14 - 5/20 (9,985)
2005	4/30 - 5/06 (7,494)
2006	6/18 - 6/24 (10,440)
2007	6/18 - 6/24 (5,483)
2008	5/28 - 6/03 (5,533)
2009	5/21 - 5/27 (7,855)
2010	4/30 - 5/06 (4,934)
2011	6/25 - 7/01 (3,647)

2+ Steelhead trout

The population estimate (or production) of 2+ steelhead trout emigrating past the trap site in lower Redwood Creek in YR 2011 equaled 5,587 individuals with a 95% CI of 4,094 – 7,081 (Figure 9). Population estimate error (or uncertainty) equaled $\pm 26.7\%$. Population abundance in YR 2011 was the second lowest of record, and 41% less than abundance for the previous seven year average ($N_{\text{avg } 7\text{yr}} = 9,943$). The average in population abundance over the eight year period equaled 9,399 (SD = 5,367; SEM = 1,898).

Correlation of time (study year) on yearly population abundances indicated a significant, negative relationship ($n = 8$, $p = 0.02$, $r = 0.79$, power = 0.74) (Figure 9). On average, there were 1,721 less individuals each study year (Figure 9). The test of time and bedload mobilizing flows (flood flows) in upper Redwood Cr (O’Kane gaging station) on population abundance also showed a significant, negative relationship over years (Regression, $p = 0.04$, adj. $r = 0.78$, negative slope for both ‘x’ variables, power = 0.37).




Figure 9. 2+ steelhead trout population abundance estimates (error bars are 95% confidence interval) in YRS 2004 – 2011. Numeric values next to box represent number of individuals. Line of best fit is a regression line, with corresponding equation, correlation value (r), and p value, lower Redwood Creek, Humboldt County, CA.

The pattern in monthly population abundance in YR 2011 was fairly uniform, and unlike the pattern for the previous seven year average (Figure 10). Monthly population abundance in YR 2011 peaked in June (N = 1,734 or 25% of the total), and for the previous seven year average peaked in May (N = 3,624 or 36% of the total) (Figure 10). In YR 2010, monthly emigration peaked in August (N = 3,086 or 51% of the total). The two most important months for 2+ steelhead trout population emigration were June and July (47% of total) in YR 2011, compared to May and June (72% of total) for the previous seven year average (Figure 10). In YR 2010, July and August were the two most important months, and accounted for 66% of the total population abundance.

The peak in weekly abundance occurred 6/25 – 7/01, and was later than previous study years with exception to YR 2010 (Table 7). For the eight study years, two peaks occurred in late April/early May, one peak occurred during the middle of May, one peak occurred in late May/early June, two peaks occurred in June, one peak occurred late June/early July, and one peak occurred in August (Table 7).




Figure 10. Comparison of 2+ steelhead trout population abundance by month in YR 2011 with the previous seven year average, lower Redwood Creek, Humboldt County, CA.

Table 7. Date of peak weekly 2+ steelhead trout population emigration by study year (number of individuals in parentheses), lower Redwood Creek, Humboldt County, CA.

Study Year	Date of peak in weekly out-migration (number in parentheses)
2004	4/30 - 5/06 (3,604)
2005	4/30 - 5/06 (2,232)
2006	6/18 - 6/24 (2,883)
2007	6/18 - 6/24 (3,066)
2008	5/28 - 6/03 (2,322)
2009	5/14 - 5/20 (314)
2010	8/20 - 8/26 (913)
2011	6/25 - 7/01 (1,283)

0+ Coho Salmon

The population estimate of 0+ coho salmon emigrating past the trap site in lower Redwood Creek in YR 2011 equaled 884 individuals with a 95% CI of 639 – 1,128 (Figure 11). Population estimate error (or uncertainty) equaled $\pm 28\%$. Population emigration in YR 2011 was the third highest of record, and about 1.3 times greater than the previous five year average. The average population abundance over the current six year period equaled 731 (SD = 698; SEM = 284).

Correlation of time (study year) on yearly population abundances indicated a non-significant, negative relationship (n = 6, p = 0.65, r = 0.23, power = 0.07) (Figure 11).




Figure 11. 0+ coho salmon population abundance estimates (error bars are 95% confidence intervals) in YRS 2006 - 2011. Lack of 95% CI for YRS 2009 and 2010 is due to scale of Y axis. Numeric values next to box represent number of individuals. Line of best fit is a regression line, with corresponding equation, correlation value (r), and p value, lower Redwood Creek, Humboldt County, CA.

The pattern in monthly population abundance in YR 2011 showed the delay in migration compared to the previous five year average (Figure 12). Monthly population abundance in YR 2011 peaked in July (N = 365 or 41% of the total), and for the previous five year average peaked in June (N = 271 or 38% of the total). In YR 2010, monthly emigration peaked in August (N = 5 or 50% of the total). The two most important months for 0+ coho salmon population emigration were June and July (73% of total) in YR 2011, and June and July (67% of total) for the previous five year average (Figure 12). In YR 2010, July and August were the two most important months, and accounted for 90% of the total population abundance. Weekly peaks in abundances occurred in early April, mid June, late June/early July, and late August (Table 8).




Figure 12. Comparison of 0+ coho salmon population abundance by month in YR 2011 with the previous five year average, lower Redwood Creek, Humboldt County, CA.

Table 8. Date of peak weekly 0+ coho salmon population emigration by study year (number of individuals in parentheses), lower Redwood Creek, Humboldt County, CA.

Study Year	Date of peak in weekly out-migration (number in parentheses)
2006	6/25 - 7/01 (113)
2007	6/11 - 6/17 (254)
2008	4/02 - 4/08 (304)
2009	6/25 - 7/01 (15)
2010	8/20 - 8/26 (5)
2011	6/25 - 7/01 (171)

1+ Coho Salmon

The population estimate (or production) of 1+ coho salmon emigrating past the trap site in lower Redwood Creek in YR 2011 equaled 113 individuals with a 95% CI of 16 – 209 individuals (Figure 13). Population estimate error (or uncertainty) equaled $\pm 86\%$. Population abundance in YR 2011 was the third lowest of record, and about 70% less than the previous seven year average ($N_{\text{avg } 7\text{yr}} = 378$). The average population abundance over the current eight year period equaled 345 (SD = 289; SEM = 102).

Correlation of time (study year) on yearly population estimates indicated a non-significant, negative relationship ($p = 0.81$, $r = 0.28$, power = 0.09) (Figure 13).




Figure 13. 1+ coho salmon population abundance estimates (error bars are 95% confidence interval) in YRS 2004 – 2011. Lack of 95% CI for YR 2010 is due to scale of Y axis. Numeric values next to box represent number of individuals. Line of best fit is a regression line, with corresponding equation, correlation value (r), and p value, lower Redwood Creek, Humboldt County, CA.

Monthly population emigration peaked in May (N = 102 or 90% of total) in YR 2011, and May (N = 290 or 77% of total) for the previous seven year average (Figure 14). In YR 2009, monthly emigration also peaked in May (N = 29 or 88% of total). The two most important months for 1+ coho salmon population emigration were April and May

(96% of total) in YR 2011, and April and May (92% of total) for the previous seven year average (Figure 14). In YR 2010, April and May were also the two most important months, and accounted for 94% of total population abundance.




Figure 14. Comparison of 1+ coho salmon population abundance by month in YR 2011 with the previous seven year average, lower Redwood Creek, Humboldt County, CA.

The peak in weekly abundance in YR 2011 occurred 5/07 – 5/13 (Table 9). For the eight study years, two peaks occurred in late April/early May, two peaks occurred in early-mid May, one peak occurred in mid May, two peaks occurred in late May, and one peak occurred in early-mid June (Table 9).

Population emigration ended during the week of 6/18 – 6/24 in YR 2011, 6/11 – 6/17 in YR 2010, 6/11 – 6/17 in YR 2009, 6/25 – 7/1 in YR 2008, 6/11 – 6/17 in YR 2007, 6/25 – 7/01 in YR 2006, 5/28 – 6/3 in YR 2005, and 6/4 – 6/10 in YR 2004.

Table 9. Date of peak weekly 1+ coho salmon population emigration by study year (number of individuals in parentheses), lower Redwood Creek, Humboldt County, CA.

Study Year	Date of peak in weekly out-migration (number in parentheses)
2004	4/30 - 5/06 (182)
2005	5/07 - 5/13 (80)
2006	6/04 - 6/10 (135)
2007	5/21 - 5/27 (32)
2008	5/21 - 5/27 (398)
2009	5/14 - 5/20 (217)
2010	4/30 - 5/06 (12)
2011	5/07 - 5/13 (85)

Cutthroat Trout

The population estimate of cutthroat trout emigrating past the trap site in lower Redwood Creek in YR 2011 equaled 118 individuals with a 95% CI of 73 – 164 individuals (Figure 15). Population estimate error (or uncertainty) equaled $\pm 39\%$. Population abundance in YR 2011 was the second highest of record. The average population abundance in YRS 2006 – 2008, and 2010 - 2011 equaled 122 (SD = 78; SEM = 35). Correlation of time (study year) on yearly population estimates indicated a non-significant, positive relationship ($p = 0.36$, $r = 0.53$, power = 0.12) (Figure 15).

Monthly population emigration peaked in August (N = 76 or 64% of total) in YR 2011, compared to July (N = 51 or 42% of total) for the average of YRS 2006 – 2008, and 2010 (Figure 16). The two most important months were July and August in YR 2011 (79% of total) and July and August for the average of YRS 2006 - 2008 and 2010 (68% of total) (Figure 16). In YR 2010, August and September were the two most important months (68% of total).

The peak in weekly abundance in YR 2011 occurred 8/13 – 8/19, and was much later than previous study years with exception to YR 2010 (Table 10). For the five study years when population abundances were determined, one peak occurred in late June/early July, two peaks occurred in July, one peak occurred in August, and one peak occurred in September (Table 10).




Figure 15. Cutthroat trout population abundance estimates in YRS 2006 – 2008 and YRS 2010 - 2011 (error bars are 95% confidence intervals). Numeric values next to box represent number of individuals. Line of best fit is a regression line, with corresponding equation, correlation value (r), and p value, lower Redwood Creek, Humboldt County, CA.




Figure 16. Comparison of cutthroat trout population abundance by month in YR 2011 with the average of YRS 2006 – 2008 and 2010, lower Redwood Creek, Humboldt County, CA.

Table 10. Date of peak weekly cutthroat trout population emigration by study year (number of individuals in parentheses), lower Redwood Creek, Humboldt County, CA.

Study Year	Date of peak in weekly out-migration (number in parentheses)
2004	
2005	
2006	7/23 – 7/29 (18)
2007	7/09 – 7/15 (15)
2008	6/25 – 7/01 (20)
2009	
2010	9/03 – 9/09 (63)
2011	8/13 – 8/19 (46)

Age Composition of Juvenile Steelhead Trout

The following percentages represent maximum values for 1+ and 2+ steelhead trout because their population estimates were compared to catches of 0+ steelhead trout (i.e. the actual catches of 0+ steelhead trout are less than expected 0+ steelhead trout population migration). Far more 1+ steelhead trout migrated downstream than either 0+ or 2+ steelhead trout in YR 2011, and for all years combined (Table 11).

Using catch and population data, the ratio of 0+ steelhead trout to 1+ steelhead trout to 2+ steelhead trout equaled 2:3.7:1 in YR 2011, 0.8:4.5:1 in YR 2010, 1:23:1 in YR 2009, 4:5:1 in YR 2008, 3:3:1 in YR 2007, 3:4:1 in YR 2006, 0.2:4:1 in YR 2005, and 1:4:1 in YR 2004.

The ratio of 1+ steelhead trout to 2+ steelhead trout was 3.7:1 in YR 2011, 4.5:1 in YR 2010, 23:1 in YR 2009, 5:1 in YR 2008, 3:1 in YR 2007, and close to 4:1 in YRS 2004 - 2006.

Table 11. Comparison of 0+ steelhead trout, 1+ steelhead trout, and 2+ steelhead trout percent composition of total juvenile steelhead trout downstream migration in YR 2011 with the previous seven year average, lower Redwood Creek, Humboldt County, CA.

Study Year	Percent composition of total juvenile steelhead trout out-migration		
	0+ steelhead*	1+ steelhead	2+ steelhead
2011	27.4	57.0	15.6
Prev. 7 Yr. Avg.	23.0	63.5	13.5
All years combined	27.3	59.0	13.7

* Uses actual catches instead of population estimate.

Fork Lengths and Weights

0+ Chinook Salmon

We measured (FL mm) 3,511 and weighed (g) 2,135 0+ Chinook salmon in YR 2011 (Table 12). Average FL (64.1 mm) and Wt (3.29 g) in YR 2011 were slightly less than the previous seven year average (Table 12). The average seasonal FL over eight study years equaled 67.2 mm (SD = 5.5 mm; SEM = 1.9 mm), and for Wt equaled 3.67 g (SD = 0.92 g mm; SEM = 0.33 g).

1+ Chinook Salmon

We measured 11 1+ Chinook salmon for FL (mm) and 10 1+ Chinook salmon for Wt (g) in YR 2011 (Table 13). Average FL (114.5 mm) and Wt (17.64 g) in YR 2011 were slightly greater than the average of years 2005, 2008 - 2010 (Table 13). The average seasonal FL using all year's data equaled 114.3 mm (SD = 6.7 mm; SEM = 3.0 mm), and for Wt equaled 16.62 g (SD = 3.40 g mm; SEM = 1.52 g).

Table 12. 0+ Chinook salmon average and median fork length (mm) and weight (g) in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.

0+ Chinook Salmon							
YR	(N)	Fork Length (mm)			Weight (g)		
		n	Avg.	Median	n	Avg.	Median
2004	554,890	3,192	59.8	61.0	1,429	2.55	2.40
2005	131,164	2,723	74.3	80.0	1,284	5.17	5.60
2006	85,149	2,058	76.2	78.0	1,715	4.96	5.10
2007	141,059	2,666	66.6	70.0	2,031	3.28	3.20
2008	173,758	3,113	64.5	67.0	2,099	3.04	3.10
2009	208,819	3,294	64.9	66.0	2,159	3.34	3.10
2010	132,733	3,543	66.8	71.0	2,125	3.71	3.50
Avg.			67.6			3.72	
2011	147,719	3,511	64.1	69.0	2,135	3.29	3.30

Table 13. 1+ Chinook salmon average and median fork length (mm) and weight (g) in YRS 2005, 2008 - 2011, lower Redwood Creek, Humboldt County, CA.

1+ Chinook Salmon							
YR	N	Fork Length (mm)			Weight (g)		
		n	Avg.	Median	n	Avg.	Median
2004	>2	2	-	-	-	-	-
2005	>11	11	109.2	111.0	11	13.60	13.50
2006	0	-	-	-	-	-	-
2007	0	-	-	-	-	-	-
2008	>10	10	113.4	113.0	9	15.80	14.2
2009*	310	57	108.8	110.0	57	14.08	14.0
2010*	22	10	125.4	126.0	10	22.00	21.9
Avg.			114.2			16.37	
2011*	15	11	114.5	117	10	17.64	19.25

* Denotes year when population abundance was determined.

0+ Steelhead Trout

We measured (FL mm) 2,354 0+ steelhead trout in YR 2011 (Table 14). Average FL (57.5 mm) in YR 2011 was greater than previous study years (Table 14). The average FL using all year's data equaled 54.3 mm (SD = 2.9 mm; SEM = 1.0 mm).

Table 14. 0+ steelhead trout average and median fork length in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.

YR	(N)	0+ Steelhead Trout*					
		Fork Length (mm)			Weight (g)		
		n	Avg.	Median	n	Avg.	Median
2004	> 18,642	2,939	49.6	52.0	-	-	-
2005	> 1,345	1,099	51.1	53.5	-	-	-
2006	> 29,957	2,757	55.8	58.0	-	-	-
2007	> 42,827	3,355	53.8	56.0	-	-	-
2008	> 39,892	2,787	52.9	56.0	-	-	-
2009	> 2,489	1,557	56.7	60.0	-	-	-
2010	> 4,566	2,275	56.9	63.0	-	-	-
Avg.			53.8		-	-	-
2011	> 9,864	2,354	57.5	58.0			

* Includes a small, but unknown number of cutthroat trout.

1+ Steelhead Trout

We measured (FL mm) 1,945 and weighed (g) 1,168 1+ steelhead trout in YR 2011 (Table 15). Average FL and Wt in YR 2011 were greater than previous study years (Table 15). The average seasonal FL over eight study years equaled 88.8 mm (SD = 3.0 mm; SEM = 1.0 mm), and for Wt equaled 8.09 g (SD = 0.89 g mm; SEM = 0.31 g).

2+ Steelhead Trout

We measured (FL mm) 653 and weighed (g) 638 2+ steelhead trout in YR 2011 (Table 16). Average FL (146.6 mm) and Wt (34.49 g) in YR 2011 were greater than previous study years (Table 16). The average seasonal FL over eight study years equaled 142.2 mm (SD = 2.2 mm; SEM = 0.8 mm), and for Wt equaled 30.88 g (SD = 1.86 g mm; SEM = 0.66 g).

Table 15. 1+ steelhead trout average and median fork length (mm) and weight (g) in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.

1+ Steelhead Trout							
YR	(N)	Fork Length (mm)			Weight (g)		
		n	Avg.	Median	n	Avg.	Median
2004	77,221	2,713	84.4	81.0	1,201	7.04	5.80
2005	32,901	1,442	90.8	89.0	919	8.31	7.40
2006	44,937	2,449	87.0	84.0	2,150	7.73	6.50
2007	37,683	2,761	88.6	87.0	2,146	7.88	7.00
2008	42,068	2,875	87.0	85.0	2,025	7.48	6.60
2009	40,728	2,349	87.6	86.0	1,673	7.59	6.80
2010	27,071	2,315	92.1	91.0	1,505	8.94	8.00
Avg.			88.2			7.85	
2011	20,501	1,945	93.2	92.0	1,168	9.77	8.85

Table 16. 2+ steelhead trout average and median fork length (mm) and weight (g) in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.

2+ Steelhead Trout							
YR	(N)	Fork Length (mm)			Weight (g)		
		n	Avg.	Median	n	Avg.	Median
2004	19,353	886	141.9	135.0	864	30.69	26.00
2005	8,754	413	143.2	139.0	412	31.25	27.05
2006	12,091	1,056	139.1	133.0	1,020	28.49	24.70
2007	12,607	1,148	141.7	134.0	1,098	31.15	25.60
2008	9,021	1,134	142.6	132.0	1,099	31.27	24.30
2009	1,743	234	140.4	133.0	219	28.65	23.90
2010	6,033	589	141.8	139.0	581	31.02	28.20
Avg.			141.5			30.36	
2011	5,587	653	146.6	145.0	638	34.49	31.65

0+ Coho Salmon

We measured (FL mm) 230 and weighed (g) 227 0+ coho salmon in YR 2011 (Table 17). Average FL and Wt in YR 2011 were the second highest of record, and similar to the previous seven year average (Table 17). The average seasonal FL over eight study years equaled 67.5 mm (SD = 7.3 mm; SEM = 2.6 mm), and for Wt equaled 4.04 g (SD = 1.33 g mm; SEM = 0.47 g).

Table 17. 0+ coho salmon average and median fork length (mm) and weight (g) in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.

YR	(N)	0+ Coho Salmon					
		Fork Length (mm)			Weight (g)		
		n	Avg.	Median	n	Avg.	Median
2004	> 202	202	66.2	66.0	198	3.76	3.50
2005	> 53	53	61.8	63.0	50	3.38	3.15
2006*	508	106	64.6	67.0	106	3.40	3.50
2007*	1,057	290	67.4	67.0	276	3.83	3.60
2008*	1,886	391	61.1	64.0	383	3.04	3.00
2009*	63	32	66.2	68.0	32	3.48	3.50
2010*	10	6	84.5	84.0	6	7.20	6.90
Avg.			67.4			4.01	
2011*	884	230	68.5	68.0	227	4.27	3.90

* Denotes study year when population abundance was determined.

1+ Coho Salmon

We measured (FL mm) 24 and weighed (g) 23 1+ coho salmon in YR 2011 (Table 18). Average FL and Wt in YR 2011 were similar to previous study years, and the previous seven year average (Table 18). The average seasonal FL over eight study years equaled 106.5 mm (SD = 3.4 mm; SEM = 1.2 mm), and for Wt equaled 13.07 g (SD = 1.20 g mm; SEM = 0.43 g).

Cutthroat Trout

We measured 57 (FL mm) and weighed (g) 54 cutthroat trout in YR 2011 (Table 19). Average FL and Wt in YR 2011 were greater than average (Table 19). The average seasonal FL over eight study years equaled 196.2 mm (SD = 17.5 mm; SEM = 6.2 mm), and for Wt equaled 80.91 g (SD = 14.24 g mm; SEM = 5.03 g).

Table 18. 1+ coho salmon average and median fork length (mm) and weight (g) in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.

1+ Coho Salmon							
YR	(N)	Fork Length (mm)			Weight (g)		
		n	Avg.	Median	n	Avg.	Median
2004	535	69	105.3	105.0	67	13.09	12.09
2005	183	39	109.4	110.0	39	13.71	13.40
2006	427	69	105.7	105.0	69	12.77	12.50
2007	102	34	104.9	107.0	34	12.36	12.30
2008	879	242	109.1	110.0	229	13.73	13.70
2009	482	101	100.0	100.0	100	10.70	10.40
2010	33	11	111.0	111.0	11	14.80	14.70
Avg.			106.5			13.02	
2011	113	24	106.7	106.5	23	13.40	12.90

Table 19. Cutthroat trout average and median fork length (mm) and weight (g) in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.

Cutthroat Trout							
YR	(N)	Fork Length (mm)			Weight (g)		
		n	Avg.	Median	n	Avg.	Median
2004	> 37	36	171.0	161.5	36	61.28	43.15
2005	> 9	9	228.7	185.0	7	70.14	64.80
2006*	97	36	193.4	182.0	35	89.80	65.60
2007*	85	44	201.7	199.0	44	97.09	84.55
2008*	54	22	178.9	163.5	21	65.87	45.10
2009	> 8	8	200.0	156.0	7	93.29	33.10
2010*	256	82	191.1	189.5	80	75.81	64.40
Avg.			195.0			79.04	
2011*	118	57	204.7	199.0	54	94.01	79.95

* Denotes study year when population abundance was determined.

Developmental Stages

1+ and 2+ Steelhead Trout

There was an obvious non-random distribution of parr, pre-smolt, and smolt designations (developmental stages) for 1+ and 2+ steelhead trout captured each study year (Table 20). A totally random distribution would equal 33.3% for each designation (parr, pre-smolt, smolt). The combined percentage of pre-smolts and smolts for 1+ steelhead trout was nearly 100%, and for 2+ steelhead trout equaled 100% (Table 20).

Table 20. Developmental stages of captured 1+ and 2+ steelhead trout in YRS 2004 - 2011, lower Redwood Creek, Humboldt County, CA.

Year	Developmental Stage (as percentage of total catch)					
	1+ Steelhead Trout			2+ Steelhead Trout		
	Parr	Pre-smolt	Smolt	Parr	Pre-smolt	Smolt
2004	0.2	31.5	68.3	0.0	5.7	94.3
2005	0.2	13.6	86.2	0.0	1.7	98.3
2006	0.1	25.1	74.8	0.0	2.1	97.9
2007	0.5	22.4	77.1	0.0	6.1	93.9
2008	0.6	15.3	84.1	0.0	1.5	98.5
2009	0.0	12.1	87.9	0.0	0.8	99.2
2010	0.1	11.1	88.8	0.0	0.5	99.5
Avg.	0.2	18.7	81.0	0.0	2.6	97.4
2011	0.1	17.0	82.9	0.0	0.7	99.3

1+ Chinook Salmon, 1+ Coho Salmon, and Cutthroat Trout

All 1+ Chinook salmon, 1+ Coho salmon, and cutthroat trout smolts captured in YR 2011 were in a smolt stage.

Trapping Mortality

The mortality of fish that were captured in the trap and subsequently handled was closely monitored over the course of each trapping period. The trap mortality (includes handling mortality) for a given species at age in YR 2011 ranged from 0.00 – 0.30%, and using all data (pooling) equaled 0.21% of the total captured and handled (Table 21). Mortality in

YR 2011 was less than average (Table 22). The variation in mortality among study years was primarily due to differences in debris loading in the trap's livebox, and whether or not large sticks/logs jammed the trap's cone.

Table 21. Trapping mortality for juvenile salmonids captured in YR 2011, lower Redwood Creek, Humboldt County, CA.

Age/spp.	Trapping Mortality in YR 2011		
	No. captured*	No. of mortalities	Percent mortality
0+ Chinook	56,706	115	0.20
1+ Chinook	11	0	0.00
0+ Steelhead	9,855	30	0.30
1+ Steelhead	2,622	1	0.04
2+ Steelhead	691	2	0.29
Cutthroat Trout	52	0	0.00
0+ Coho	226	0	0.00
1+ Coho	24	0	0.00
Overall:	70,187	148	0.21

* Not expanded for missed day(s) catch during periods of trap non-deployment.

Table 22. Comparison of trapping mortality of juvenile salmonids in eight consecutive study years, lower Redwood Creek, Humboldt County, CA.

Study Year	Trapping Mortality		
	No. captured*	No. of mortalities	Percent mortality
2004	88,088	167	0.19
2005	14,736	143	0.97
2006	55,671	93	0.17
2007	92,165	57	0.06
2008	126,201	140	0.11
2009	59,655	1,064	1.78
2010	52,649	113	0.21
2011	70,187	148	0.21
Avg.	69,919	241	0.46
Pooled	559,352	1,925	0.34

* Not expanded for missed day(s) catch during periods of trap non-deployment.

Stream Temperatures

The average daily (24 hr period) stream temperature from 4/09/11 – 9/09/11 was 14.5 °C (or 58.1 °F) (95% CI = 13.9 – 15.0 °C), with daily averages ranging from 8.1 – 18.6 °C (46.6 – 65.5 °F). Median daily temperature in YR 2011 equaled 16.1 °C (or 61.0 °F). Average stream temperatures during the trapping periods in YRS 2004 – 2011 were similar, with the largest difference among years equaling 1.4 °C (Table 23). However, average daily stream temperature (during the trapping period) over study years significantly decreased over time (Correlation, $p = 0.01$, $r = 0.83$, slope is negative, power = 0.85, alpha = 0.10). Similar to past data, the average daily stream temperatures during the trapping period in YR 2011 were inversely related to the average daily stream discharge during the trapping period (Regression, $p < 0.000001$, $R^2 = 0.84$, slope is negative, power = 1.0). The minimum stream temperature in YR 2011 equaled 7.4 °C (45.3 °F) and occurred on 4/09/11; the maximum stream temperature equaled 21.8 °C (71.2 °F) and occurred on 7/28/11 (Table 23).

Table 23. Average, minimum, and maximum stream temperatures (°C) (standard error of mean in parentheses) at the trap site during the trapping periods in YRS 2004 – 2011, lower Redwood Creek, Humboldt County, CA.

Study Year	Stream Temperature					
	Celsius			Fahrenheit		
	Avg.	Min.	Max.	Avg.	Min.	Max.
2004	15.5 (0.2)	9.3	22.6	60.0 (0.8)	48.7	72.3
2005	15.6 (0.3)	9.0	22.6	60.1 (0.5)	48.2	72.3
2006	15.5 (0.3)	7.1	23.1	60.0 (0.5)	44.8	73.6
2007	15.3 (0.3)	8.1	24.2	59.5 (0.6)	46.6	75.6
2008	14.2 (0.3)	6.9	21.8	57.6 (0.5)	44.4	71.2
2009	14.7 (0.2)	7.4	21.2	58.5 (0.4)	45.3	70.2
2010	14.6 (0.2)	8.2	21.3	58.3 (0.4)	46.8	70.3
2011	14.5 (0.3)	7.4	21.8	58.0 (0.5)	45.3	71.2
Avg.	15.0 (0.2)			59.0 (0.4)		

Average monthly stream temperatures during the majority of the trapping season (April – July) in YR 2011 ranged from 9.2 – 17.4 °C (48.6 – 63.3 °F) (Table 24). Highest stream temperatures occurred in the later part of the trapping season (June and July) each study year. The MWAT during the trapping period in YR 2011 at the trap site was 18.1 °C (64.6 °F); and occurred on 7/29/11 (Table 25). MWMT in YR 2011 was 21.2 °C (70.2 °F) and occurred on 7/29/11 as well (Table 25). MWAT and MWMT in YR 2011 were similar to previous study years (Table 25).

The average daily stream temperature (transformed) increased over the study period (days, transformed) in YR 2011 (Correlation, $p < 0.000001$, $r = 0.90$, slope is positive, power = 1.0) (Figure 16), as well as in past study years (Figure 17).

Table 24. Average monthly stream temperature (°C) (°F in parentheses) at the trapping site in study years 2004 - 2011, lower Redwood Creek, Humboldt County, CA.

Study Year	Average stream temperature in Celsius (°F in parentheses)				
	April	May	June	July	Avg.
2004	11.9 (53.4)	14.7 (58.5)	16.8 (62.2)	18.6 (65.5)	15.5 (59.9)
2005	11.5 (52.7)	12.8 (55.0)	14.6 (58.3)	18.5 (65.3)	14.3 (57.7)
2006	10.4 (50.7)	13.9 (57.0)	16.7 (62.1)	18.2 (64.8)	14.8 (58.6)
2007	10.7 (51.3)	13.4 (56.1)	16.4 (61.5)	18.5 (65.3)	14.8 (58.6)
2008	9.8 (49.6)	13.5 (56.3)	15.6 (60.1)	17.4 (63.3)	14.1 (57.4)
2009	10.7 (51.3)	12.9 (55.2)	15.7 (60.3)	17.3 (63.1)	14.2 (57.6)
2010	10.1 (50.2)	11.1 (52.0)	13.8 (56.8)	17.2 (63.0)	13.1 (55.6)
2011	9.2 (48.6)	11.3 (52.3)	14.5 (58.1)	17.4 (63.3)	13.1 (55.6)
Avg.	10.5 (50.9)	13.0 (55.4)	15.5 (59.9)	17.9 (64.2)	

Table 25. Maximum weekly average temperature (MWAT) and maximum weekly maximum temperature (MWMT) for stream temperatures °C (°F in parentheses) at the trap site in lower Redwood Creek, Humboldt County, CA., study years 2004 - 2011.

Study Year	MWAT		MWMT	
	Date of Occurrence	°C (°F)	Date of Occurrence	°C (°F)
2004	7/22/04	19.3 (66.7)	7/18/04	22.2 (72.0)
2005	7/17/05	19.2 (66.6)	7/17/05	22.1 (71.8)
2006	7/25/06	19.2 (66.6)	7/25/06	22.7 (72.9)
2007	7/21/07	19.2 (66.6)	7/31/07	22.4 (72.3)
2008	7/07/08	18.2 (64.8)	7/07/08	21.1 (69.8)
2009	8/11/09	18.3 (64.9)	8/11/09	20.9 (69.7)
2010	7/14/10	17.9 (64.3)	7/17/10	20.8 (69.4)
2011	7/29/11	18.1 (64.6)	7/29/11	21.2 (70.2)




Figure 17. Average, minimum, and maximum stream temperature (°C) in lower Redwood Creek, Humboldt County, CA., 2011.




Figure 18. Average daily stream temperature (°C) in YRS 2004 – 2011, lower Redwood Creek, Humboldt County, CA.

DISCUSSION

The main goal of our downstream migration study in lower Redwood Creek is to estimate and monitor the production of Chinook salmon, steelhead trout, coho salmon, and cutthroat trout from the majority of the Redwood Creek watershed in a reliable, long-term manner. The long term goal is to monitor trends in smolt abundance and smolt size, and to detect positive or negative changes due to watershed conditions and restoration activities in the basin. Redwood Creek is a difficult stream to monitor adult salmon and steelhead populations on a long term basis using traditional techniques (weirs and spawning ground surveys) due to adult salmon and steelhead run timing, water depth, precipitation, hydrology, and stream turbidity. However, “quantifying juvenile anadromous salmonid populations as they migrate seaward is the most direct assessment of stock performance in freshwater” (Seiler et al. 2004). In addition, studies in various streams have found that smolt numbers can relate to stream habitat quality, watershed condition, restoration activities, the number of parents that produced the cohort, and future adult populations.

The eighth consecutive year of trapping in lower Redwood Creek occurred during a WY with a slightly greater stream flow compared to the historic value, and the average monthly stream discharge during the majority of the trapping season (April – July) in YR 2011 was 1.4 times greater than the historic average. The environmental conditions for downstream migrant trapping in YR 2011 were more difficult to operate the trap in compared to most study years, and we missed a total of five days of trapping due to high flow events and potentially, high debris loading in the trap’s livebox. However, the estimates for catch and subsequent expansions to the population level, based on the missed trapping days, were negligible for each species at age; the greatest impact on a population estimate was estimated at 4%, and the adjusted point value easily fell within the 95% confidence interval of the un-adjusted point estimate. The number of fish missed when the trap was inoperable would not have greatly impacted population estimates. Thus, this season’s trapping resulted in very good estimates of wild Chinook salmon, steelhead trout, cutthroat trout, and coho salmon smolt abundances from areas upstream of the trapping site. The abundance estimate for 1+ coho salmon smolts (+ or – 86%) was not as precise as for other species at age, yet sample sizes for marking these fish were much lower. The decrease in precision should not preclude the fact that yearling coho salmon smolts were in low abundance in YR 2011 ($N < 114$ individuals).

0+ Chinook Salmon

0+ Chinook salmon (ocean-type) were the most numerous migrant captured in seven of eight consecutive study years. The correlation of 0+ Chinook salmon trap catches, study years, and whether or not there were flood type flows in the upper basin during egg incubation/embryogenesis with study years was significantly negative ($p = 0.02$). However, the best model to describe the variation in trap catches over years included the single variable of flood type flows in the upper basin ($p = 0.0004$). Inference from these models suggest that the trap catches in lower Redwood Creek were strongly tied to

Chinook salmon in the mid to upper basin, and to environmental conditions in the mid to upper basin with respect to stream flow, redd formation, and survival to emergence from redds.

On a population basis, 0+ Chinook salmon smolts were 1.5 – 5.7 times more abundant than 1+ and 2+ steelhead trout smolts combined over the eight years of study. The population abundance of 0+ Chinook salmon in YR 2011 ($N = 147,719$) was the fifth lowest of record, 1.11 times greater than abundance in YR 2010, 74% less than the highest abundance measured in YR 2004 ($N = 554,890$), and 28% less than abundance for the average of the previous seven years. The overall trend in abundance over eight consecutive study years was negative; however, statistical significance was not detected, most likely due to the observed variation in population abundances and low sample sizes ($n = 8$ data points). Testing trends in abundance often requires numerous years of data to determine a statistically, reliable trend. Trends with low sample sizes not only preclude statistical significance, but the slope of the trend line can also change with the addition or omission of a single data point. For example, the slope of the trend line in Chinook salmon smolt abundance over eight years was greatly influenced by the high abundance determined in YR 2004; if we removed YR 2004 data, the slope would be positive instead of negative. The abundance in YR 2004 could be considered an outlier, however, we did catch more fish in that year than all years except for YR 2008, even though the stream gradient in YR 2004 was much less than the gradient in YRS 2006 - 2010. At this trapping site, trapping efficiencies are positively related to the stream gradient. In addition, population abundance in upper Redwood Creek peaked at about 630,000 individuals in YR 2004, which was much higher than other study years in that data set. Thus, I did not consider YR 2004 data in lower Redwood Creek to be unreasonable, or an outlier. Based upon data collected in upper Redwood Creek, it may take nine plus years to determine a significant trend in 0+ Chinook salmon population abundance passing through lower Redwood Creek (Sparkman 2011b). Although the trend in abundance in lower Redwood Creek was non-significant, there was a significant, negative relationship in abundance when flood type flows in the upper basin were added as a dummy variable to the regression model ($p < 0.10$). This did not come as a surprise because there is a direct relationship between population abundances in the upper basin with that of the lower basin, and population abundances in the upper basin are strongly affected by flood type flows during adult spawning and the redd incubation period (Sparkman 2011b).

The overall reduction in population abundance we observed from YRS 2004 – 2011 in lower Redwood Creek could be due to: 1) decrease in the total number of spawners upstream of the trap site, 2) high bedload mobilizing flows during and after reproduction which scoured or jostled redd gravels in the mid to upper basin, or 3) some combination of factors 1 and 2. Changes in spawner distributions within the Redwood Creek basin are not likely responsible for the decrease because Chinook salmon do not generally spawn in mainstem areas below the trap site in lower Redwood Creek, and the number of spawners in Prairie Creek in YR 2011 was near average (Duffy, pers. comm. 2011). Currently, we cannot separate effects of lower adult population abundance during years with high, bedload mobilizing flows on the subsequent production of juveniles because: 1) adult counts are only recently being conducted, 2) peak flows capable of redd scour in

lower Redwood Creek occurred each study year (YRS 2004 – 11), and 3) peak flows capable of redd scour in mid to upper Redwood Creek occurred in YRS 2005 and 2006. Several investigators have shown that the scour of redds due to high streamflows or floods can often cause severe decreases in the production of juvenile salmonids (Gangmark and Bakkala 1960, McNeil 1966, Holtby and Healey 1986, Montgomery et al. 1996, Devries 1997, Schuett-Hames et al. 2000, Seiler et al. 2003, Don Chapman pers. comm. 2003, Greene et al. 2005); and that estimates of mortality attributable to high flows and redd scour can reach 90% (Schuett-Hames et al. 2000). Greene et al. (2005) were able to show that the flood recurrence interval (and magnitude of floods) during Chinook salmon intragravel development was the second most important variable in their models used to predict the return rate of adult Chinook salmon. They further report that “large flow events may be a key factor in regulating Chinook salmon populations in the Skagit River basin, Washington” (Greene et al. 2005). Since each study year peaks in streamflow (11,000⁺ cfs) capable of redd scour occurred in lower Redwood Creek with exception to YR 2010, and no peaks capable of redd scour occurred in the upper basin in YR 2011, the most plausible reason for the decrease in abundance in YR 2011 is a simple decrease in the number of returning adults. Although adult harvest is not allowed in Redwood Creek, the ocean salmon sport fishery does harvest an unknown number of adult Chinook salmon that belong to Redwood Creek Roper and Scarnecchia (1999) reported that the abundance of 0+ Chinook salmon in the South Umpqua River, Oregon was positively related to the number of returning adults.

Subsequent to winter flows and emergence from redds, 0+ Chinook salmon migrated downstream nearly each day during the trapping period in YR 2011. Weekly peaks in abundance during a given study year were relatively large, ranging from 27,057 – 111,000 individuals. In YR 2011 the peak in weekly abundance equaled about 27,000 individuals, and occurred in late June/early July. The pattern in population abundance by month in YR 2011 compared to the previous seven year average showed the temporal delay in migration in YR 2011 (similar to YR 2010). Peak monthly emigration occurred in July in YR 2011, compared to June for the previous seven year average. The two most important months in YR 2011 were June and July which was also the two most important months for the previous seven year average. The delay in peak monthly migration in YR 2011 was most likely attributable to higher than average stream flows during the trapping period. Population emigration during a given month can be quite large, reaching values up to 292,000 (June 2004), and in June 2011 population abundance equaled 75,872 individuals.

Each study year 0+ Chinook salmon (ocean-type) emigrating from Redwood Creek exhibit two different juvenile life histories (fry and fingerling) based on size and time of downstream migration. The fry (Avg. FL = 40 mm in YR 2011) are migrating shortly after emergence from spawning redds, and therefore are much smaller than the fingerlings (or smolts) (Avg. FL = 71 mm in YR 2011) which have reared in the stream for a longer period of time prior to passing the trap site. Although there is overlap in downstream migration, temporal differences are evident by two peaks in migration when data for fry abundance and fingerling abundance are plotted using two ‘y’ axes (Sparkman 2011a). For example, the first weekly peak (albeit relatively small, N =

12,564) in population emigration in YR 2011 occurred during 4/23 – 4/29, and consisted of 97% fry and 3% fingerlings. The average size of fry during this time period equaled 40 mm. Weeks later, the largest peak in abundance (6/25 – 7/01, N = 27,057) consisted solely of fingerlings with an average FL of 67 mm. Fry and fingerlings that are marked for efficiency trials in the upper basin are captured each year by the trap in lower Redwood Creek. Factors that can influence the temporal component to fry and fingerling migration are: 1) time of adult spawning, 2) how far upstream of the trap site the adults spawned, 3) time from egg deposition to fry emergence from redds, and 4) travel rate, among other factors.

Small numbers of fry relative to the number of fingerlings migrated downstream through lower Redwood Creek each study year, with exception to YRS 2004 and 2011. The greatest number of fry migrating downstream towards the estuary and ocean occurred in YR 2004 (N = 82,584), and in YR 2011, 19% (or 27,809 individuals) of the population consisted of fry. The percentage of fry in the 0+ Chinook salmon population over the previous seven years ranged from 0.1 – 15%, and averaged 5%. The average number of fry over the previous seven years equaled about 8% of the Chinook population. 0+ Chinook salmon fingerlings comprised the majority of the population each year, with percentages ranging from 81 – 99.9% of the total abundance; the greatest number of fingerlings (N = 472,306) migrating downstream occurred in YR 2004 as well. In YR 2011, 81% (or 119,910) of the 0+ Chinook salmon population consisted of fingerlings. In contrast, the 0+ Chinook salmon population emigrating from upper Redwood Creek consisted of nearly equal numbers of fry (113,254 or 51%) and fingerlings (108,475 or 49%) when averaged over a twelve year period (YRS 2000 – 11), and in YR 2011, 62% of the 0+ Chinook salmon emigrant population passing through upper Redwood Creek consisted of fry (Sparkman, In progress). Clearly areas upstream of the trap site in upper Redwood Creek are important for adult Chinook salmon spawning; and by the time the juvenile Chinook salmon population are passing through lower Redwood Creek, most are in the fingerling (smolt) size category.

The average size (FL = 64 mm; Wt = 3.3 g) of 0+ Chinook salmon emigrants in YR 2011 was the second lowest of record, and less than the previous seven year average.

1+ Chinook Salmon

One year old juvenile Chinook salmon (stream-type) in Redwood Creek represent the third juvenile Chinook salmon life history. Stream-type juvenile Chinook salmon are easily differentiated from ocean-type by size at time of downstream migration, and general appearance. The average size (FL mm) in April/May, 2011 for example, was 109 mm for 1+ Chinook salmon and 41 mm for 0+ Chinook salmon.

1+ Chinook salmon in Redwood Creek appear to be in very low abundance as evidenced by trap catches totaling 104 individuals over eight consecutive study years. In YR 2011 we captured 11 1+ Chinook salmon, and for the third consecutive year determined a mark/recapture population estimate (N = 15 in YR 2011). The population estimate was

310 in YR 2009 and 22 in YR 2010. The majority of 1+ Chinook salmon migrated downstream during May in YRS 2009 - 2011.

When present, 1+ Chinook salmon in Redwood Creek are more likely to be progeny of fall/winter-run Chinook salmon adults than from spring-run adults because few if any spring-run Chinook salmon are observed during spring and summer snorkel surveys in Redwood Creek (Dave Anderson, pers. comm. 2011). For example, in 24⁺ years of adult summer steelhead snorkel dives, adult spring Chinook salmon were only observed in one year (1988) and in very low numbers (< 7 individuals) (Dave Anderson, pers. comm. 2011). Additionally, streamflows during late spring/summer months can become so low that adult upstream passage into upper Redwood Creek can become problematic. High average stream temperatures (eg > 20 °C) may also prevent any adult spring-run Chinook salmon migration into upper Redwood Creek, or inhibit their ability to over-summer in pools. The sedimentation in Redwood Creek and subsequent decrease in pool depths and other stream habitat may also limit the ability to over-summer as well.

Thus, a spring run of Chinook salmon adults was probably not responsible for the production of yearling Chinook salmon juveniles in Redwood Creek. Bendock (1995) also found both stream-type and ocean-type juvenile Chinook salmon in an Alaskan stream which only has one adult Chinook salmon race; and Conner et al. (2005) reported that fall Chinook salmon in the Snake River produced juveniles exhibiting an ocean-type or stream-type juvenile life history. Teel et al. (2000) found that for some populations of coastal Chinook salmon, ocean-type and stream-type juveniles were genetically undifferentiated, and probably arose from a common ancestor. They further conclude that the stream-type life history probably evolved after the ocean-type colonized (post glacial period) the rivers in study. An important question which may be unanswerable, is whether the one year old life history for juvenile Chinook salmon in Redwood Creek was more prevalent prior to the changes in the watershed associated with land use activities, flood events, and natural geologic processes. Perhaps with continued stream restoration and maturation of the riparian zone in Redwood Creek, we will see an increase in the numbers of stream-type Chinook salmon smolts. The 1+ Chinook salmon life history pattern may be important for increased ocean survival of Chinook salmon juveniles, and general species diversity (author, Don Chapman pers. comm. 2003).

0+ Steelhead Trout

The number of 0+ steelhead trout that can remain upstream of the trap site is considered to be some function of a fish's disposition to out-migrate (or not out-migrate) and habitat carrying capacity. Meehan and Bjornn (1991) comment that juvenile steelhead trout have a variety of migration patterns that can vary with local conditions, and that the trigger for out-migration can be genetic or environmental. They further state that some steelhead populations normally out-migrate soon after emergence from redds to occupy other rearing areas (we observe this as well in both upper and lower Redwood Creek). Habitat carrying capacity is generally thought to be related to environmental (hydrology, geomorphology, stream depth and discharge, stream temperatures, cover, sedimentation,

etc.) and biological variables (food availability, predation, salmonid behavior), and any interactions between the two (Murphy and Meehan 1991). The general idea is that when habitat carrying capacity is exceeded (e.g. over-seeding, surplus production), the juvenile fish emigrate to find other areas to rear. A problem with the view of habitat carrying capacity's affect on migration is that it fails to explain why juvenile salmonids (e.g. 0+ SH, 0+ CO) emigrate at low, upstream densities or low, upstream population levels.

Young-of-year steelhead trout downstream migration in Redwood Creek is considered to be stream redistribution (passive and active) because juvenile steelhead in California normally smolt and enter the ocean at one to two years old, with lesser numbers out-migrating at an age of 3⁺ years (Busby et al. 1996, Sparkman 2011b). Perhaps the most important finding with respect to 0+ steelhead trout movements in Redwood Creek were from experiments conducted in YRS 2006, 2007, 2010, and 2011 in which we marked 0+ steelhead trout (FL 40 – 60 mm) in the upper basin to later see if we would recapture a given percentage at the trap in lower Redwood Creek. Recaptures occurred within each study year of the experiments, and in YR 2010 we recaptured six out of 100 marked with a upper caudal fin clip (snip). In YR 2011 we only recaptured 1 out of 100 that were marked and released at the trap in upper Redwood Creek. To the best of my knowledge, these were the first experiments to show 0+ steelhead trout may cover considerable distances (e.g. 29 mi.) while moving downstream in search of rearing areas.

Trap catches of 0+ steelhead trout in YR 2011 (n = 9,864) in lower Redwood Creek were greater than catches for 1+ steelhead trout in YR 2011, and much greater than catches of 2+ steelhead trout in YR 2011. Trap catches of 0+ steelhead trout in YRS 2004 – 2011 ranged from 1,345 – 42,827 individuals, and averaged 18,698. In comparison to population abundances of 1+ and 2+ steelhead trout, 0+ steelhead trout comprised 27% of the total juvenile steelhead trout out-migration over the current, eight year period. With respect to 2+ steelhead trout, the ratio of 0+ steelhead trout catches to 2+ steelhead trout population abundances over the eight year period ranged from 0.8:1 to 4.4:1, and averaged 1.9:1. Relatively high catches of young-of-year steelhead trout by downstream migrant traps in small and large streams is not uncommon (USFWS 2001, Rowe 2003, Johnson 2004, Don Chapman pers. comm. 2004, Sparkman 2011b). For example, 0+ steelhead trout catches in upper Redwood Creek from YRS 2000 – 2011 ranged from 32,585 - 128,885 and averaged 69,460 per year (Sparkman, In progress). Similar to 0+ Chinook salmon, mid to upper Redwood Creek is important for adult steelhead trout spawning.

The 0+ steelhead trout captured by the trap in lower Redwood Creek indicate these fish are going to rear for some time period in lower Redwood Creek (including the estuary), or Prairie Creek. Dave Anderson (pers. comm. 2011), for example, routinely captures young-of-year steelhead trout (and coho salmon) in the estuary during summer and early fall sampling. Although relatively few 0+ steelhead trout migrated downstream past the trap site in YR 2011, the condition of lower Redwood Creek and the estuary can impact 0+ steelhead trout, which in turn could influence the number of older, juvenile steelhead trout in following years.

1+ Steelhead Trout

One-year-old steelhead trout smolts were the most numerous juvenile steelhead trout migrating downstream through lower Redwood Creek in at least six of eight consecutive study years. In YR 2011, 1+ steelhead trout outnumbered 0+ and 2+ steelhead trout, with the corresponding ratio of 2.1:1 for 1+ steelhead trout to 0+ steelhead trout; and 3.7:1 for 1+ steelhead trout to 2+ steelhead trout. The ratio of 1+ steelhead trout to 0+ steelhead trout averaged 7:1 over the current eight years, and for 1+ to 2+ steelhead trout populations averaged 6.3:1. On a percentage basis, 1+ steelhead trout comprised 79 – 96% of the total juvenile steelhead trout age 1 and older population abundance each study year. In YR 2011, there were 3.7 times as many 1+ steelhead trout than 2+ steelhead trout passing through lower Redwood Creek. Obviously, 1+ steelhead trout comprise the majority of steelhead trout smolts passing the trap site.

Population abundance in YR 2011 ($N = 20,501 \pm 17\%$) was the lowest of record, and 53% less than the previous seven year average. With inclusion of YR 2011 data, the trend in abundance over the eight year period was statistically negative ($p = 0.03$), and on average, we are seeing about 5,172 less individuals each study year. The high abundance in the first year (YR 2004, $N = 77,221$) may appear to ‘drive’ the regression, however, the slope is still negative when YR 2004 data is omitted. The best model describing the trend in abundance over time included study year and whether or not there were flood type flows in the upper basin each study year ($p = 0.02$). The relationship with flood type flows was negative, and may indicate poor overwinter survival when stream flows reach a critical threshold ($cfs > 6,000$) in mid to upper Redwood Creek. Redwood Creek follows a fault for many miles, and this fault probably limits the amount and degree of channel meandering and formation of off channel refugia (alcoves, backwaters, etc.). In addition, the lack of large woody debris limits overwintering habitat.

In addition to differences in population abundance among study years, there were temporal differences in monthly emigration in YR 2011 compared to the previous seven year average. The most important month for migration was July in YR 2011, compared to June for the previous seven year average, and the two most important months for migration were June and July in YR 2011, compared to May and June for the previous seven year average. In addition, migration in YR 2011 continued into September, when normally migration is over by early to mid August. The prolonged migration observed in YR 2011 was most likely due to higher than average stream flow during the study period (similar to YR 2010), compared to historic values and previous study years (excluding YR 2010). Population emigration during a given month can be quite large, reaching values up to 33,000 (May 2004), and in July of 2011, population abundance peaked at 7,630 individuals.

The average size of 1+ steelhead trout migrants in YR 2011 (93 mm, 9.8 g) was greater than previous study years, and may be attributable to the prolonged migration and higher than average stream flows observed in YR 2011. Previous travel time and growth studies on pit tagged 1+ steelhead trout emigrating from the upper basin to be recaptured at the

trap in lower Redwood Creek showed that 1+ steelhead trout grew more as travel time increased (Sparkman 2010).

Information in the literature indicates steelhead smolting at age 1 is not uncommon, particularly in streams that are south of British Columbia (Quinn 2005, Busby et al. 1996). The percentage of 1+ steelhead trout showing parr characteristics in Redwood Creek was very low each study year (0.0 - 0.6%), and indicated that few 1+ steelhead trout migrated downstream in a stream-residence form (parr). In contrast, the majority of 1+ steelhead trout (68 – 89%) in a given study year were emigrating in a smolt stage. Given more data years, we may find relationships between developmental stages and physical variables measured in the stream. For example, I found that the percentages of 1+ steelhead trout showing smolt characteristics each year (YRS 2000-09) in upper Redwood Creek were negatively related to water temperatures ($n = 10$, $p < 0.05$) (Sparkman 2010). Quinn (2005) reported both photo period and stream temperature play important roles in smoltification by providing an external stimulus for the endocrine system, which in turn drives the internal physiological changes necessary for smoltification.

1+ steelhead trout are actively migrating from the upper basin to the lower basin as evidenced by trap catches in lower Redwood Creek of efficiency trial fish and pit tagged fish released from the upper trap site. The marked 1+ steelhead trout emigrating from upper Redwood Creek and through lower Redwood Creek have also been captured in the estuary (Dave Anderson, pers. comm. 2011) since the beginning of our smolt trapping studies. 1+ steelhead trout marked and released at the lower trap (for trap efficiencies) have also been captured in the estuary each study year (Dave Anderson, pers. comm. 2011). We have not observed re-migration of 1+ steelhead trout into lower or upper Redwood Creek based upon elastomer marked releases in YR 2001 ($n = 374$), YR 2004 ($n = 577$), and YR 2005 ($n = 146$); and pit tagged releases in YRS 2005 ($n = 46$), 2006 ($n = 246$), 2007 ($n = 484$), 2008 ($n = 203$), 2009 ($n = 417$), and 2010 ($n = 260$). All 2+ steelhead trout captured by the traps were inspected for marks and scanned for pit tags, which would have been applied at age-1. These tests confirmed that the elastomer marked and pit tagged fish did not migrate back upstream to rear for another year and emigrate as age-2 steelhead trout smolts. Elastomer mark retention was assumed to be adequate for the studies because Fitzgerald et al. (2004) assessed elastomer mark retention in Atlantic salmon smolts and found that tag retention in the lower jaw was greater than 90% for the first 16 months. Pit tag retention was also assumed to be sufficient based upon a study by Newby et al. (2007).

As previously mentioned, far more 1+ steelhead trout emigrated past the lower trap than older, juvenile steelhead trout age-classes (2+). The ratio of 1+ steelhead trout:2+ steelhead trout in YR 2011 equaled 14:1 in upper Redwood Creek, 3.7:1 in lower Redwood Creek, and 3.1:1 in Prairie Creek. The ratio of 1+ steelhead trout downstream migration is not unique to Redwood Creek, and other downstream migration studies have routinely documented 1+ steelhead trout emigration (USFWF 2001, Ward et al. 2002, Johnson 2004; B. Chesney pers. comm. 2011, among many others). However, the ratio of 1+ steelhead trout to 2+ steelhead trout (near 4:1 each study year) passing through

lower Redwood Creek was much different than that determined in a nearby river (Mad River), which equaled 1:6 in YR 2001 and 1:3 in YR 2002 (Sparkman 2002). Whether these differences are indicative of stream conditions or attributable to the different stock in each stream is unknown. In the Keogh River, about 20% of the total steelhead trout smolt yield consisted of 1+ steelhead trout parr (McCubbing and Ward 2003).

Based upon studies in other streams, the number of returning adult steelhead trout that migrated to the ocean as one-year-old smolts is relatively low, and usually less than 29% (Pautzke and Meigs 1941, Maher and Larkin 1955, Busby et al. 1996, McCubbing 2002, McCubbing and Ward 2003). Based upon a limited number of scale samples from adult steelhead trout (n = 10) collected in Redwood Creek, 30% of the adults entered the ocean as one-year-old juveniles; the most successful juvenile steelhead migrants to reach adulthood were 2+ steelhead trout (author). More recently, data collected from adults in YR 2007/2008 showed that 50% of the adults had entered the ocean as a one year old smolt, and in YR 2008/09 the percentage equaled 40%. CDFG AFRAMP is currently collecting scale samples from adult steelhead trout in Redwood Creek to increase sample size (author, in progress).

The percentage of adult steelhead trout that smolt and enter the ocean at age-1, and the reason(s) for the relative large number of 1+ steelhead trout emigrating from the basin of Redwood Creek warrants further investigation. Our pit tagging experiments with 1+ steelhead smolts should provide useful insights when conducted over multiple consecutive years because if most of the 1+ steelhead trout are not actually entering the ocean, we should then be able to recapture a given percentage of those fish the following year with the rotary screw trap in lower Redwood Creek and seine nets in the estuary; if we fail to recapture any of the marked 1+ steelhead trout the following year, then a logical conclusion would be that the fish either stayed in the stream and suffered severe mortality during winter, actually entered the ocean, or some combination of the two factors. To date, we have not recaptured any 2+ steelhead trout that were marked as 1+ steelhead trout the previous year. Thus, our data is showing, in combination with adult scale analyses, that 1+ smolts are entering the ocean at age-1.

I hypothesize that 1+ (and 0+) steelhead trout have changed their life history to limit the time spent in freshwater in order to avoid high, and at times, lethal stream temperatures that occur during summer months. In YR 2006 we observed and documented lethal stream temperatures in upper Redwood Creek, and every summer in late July we observe maximums in stream temperatures that range from 24.4 – 29.5 °C (or 75.9 – 85.1 °F) (Sparkman 2011b). In addition, stream flow during summer months is very low, which decreases the amount of space available for rearing. Over-summer conditions, particularly in mid to late July, could be limiting the production of older age classes (2+ steelhead trout) in Redwood Creek. However, we may see decreases in sedimentation and stream temperatures due to restoration activities in the basin, which in turn should provide more suitable over-summer conditions for 0+ and 1+ steelhead trout.

2+ Steelhead Trout

In several studies investigating steelhead trout life histories, the majority of the returning adult steelhead spent two or more years as juveniles in freshwater prior to ocean entry (Pautzke and Meigs 1941, Maher and Larkin 1955, Busby et al. 1996, Smith and Ward 2000, McCubbing 2002, McCubbing and Ward 2003). Pautzke and Meigs (1941), for example, reported that 84% of returning adult steelhead trout in the Green River had spent two or more years as juveniles in freshwater. Maher and Larkin (1955) found that 98% of the adult steelhead they examined had spent two or more years in freshwater prior to entering the ocean, McCubbing (2002) reported 92% of steelhead adults in a British Columbia stream had spent two or more years as juveniles in freshwater, and McCubbing and Ward (2003) reported that 71% of the adult returns in YR 2003 had entered the ocean as 2 or 3 year old smolts. If this applies to steelhead trout in Redwood Creek, then 2+ steelhead trout are the most important (and most direct) group of juvenile steelhead trout that contribute to future adult steelhead trout populations. The paradox for the 2+ steelhead trout smolt in Redwood Creek is that they were far less abundant (by about 67 - 96%) than 1+ steelhead trout smolts in any given study year. With respect to the combined population of 1+ and 2+ steelhead trout smolts each year, 2+ steelhead trout comprised 4 – 25% of the population.

The population abundance of 2+ steelhead trout in YR 2011 was the second lowest of record, and about 41% less than the average abundance over the previous seven years. The sharp decline in YR 2009 caused the previously non-significant negative trend to become a statistically significant negative trend, and with inclusion of YRS 2010 and 2011 data, the trend remained significantly negative. On average, there were 1,721 less individuals each study year. The inclusion of flood type flows in the linear model was significantly negative, however, the best model included population abundance and time (years). As discussed in the section for 0+ Chinook salmon, testing trends in abundance often requires numerous, consecutive years of data to determine a reliable trend. For example, the slope of the trend line for 2+ steelhead trout emigrating through lower Redwood Creek was greatly influenced by the high population abundance measured in YR 2004. However, if we considered YR 2004 data an outlier and removed YR 2004 data from the correlation test, we would still have a negative relationship, albeit non-significant. The negative trend in abundance for 2+ steelhead trout emigrating from upper Redwood Cr was first detected at year four, and statistically determined at year eight. The 2+ steelhead trout population emigrating from upper Redwood Creek during study years 2000 – 2011 are currently showing a significant, negative trend (Sparkman, In progress). Whether the populations of 2+ steelhead trout smolts passing through lower Redwood Creek are showing a true negative trend will take more study years to determine.

Confidence intervals (and percent error) for the population abundance estimate of 2+ steelhead trout passing through lower Redwood Creek each year were larger than the 95% confidence intervals for 1+ steelhead trout because: 1) 2+ steelhead trout are typically harder to catch than younger age-classes of steelhead trout, and 2) sample sizes for marking and subsequent recapture were lower. During the trapping period we

routinely adjusted the trap configuration and installed weir panels to increase the capture efficiency of 2+ steelhead trout. Additionally, we performed numerous mark/recapture trials in a given week, and when combined with altering trap configuration and paneling, were then able to produce a reliable population estimate with a low to moderate error term (e.g. $\pm 27\%$ observed in YR 2011).

In addition to differences in population abundance among study years, there were temporal differences in monthly emigration. Migration in YR 2011 was more spread out and rather uniform among months. In YR 2011, slightly more of the 2+ steelhead trout smolts emigrated during June (25% of total), compared to May (36% of total) for the previous seven year average. The two most important months were June and July in YR 2011, compared to May and June for the previous seven year average. The weekly peak in emigration in YR 2011 was also later than previous study years, with exception to YR 2010. Similar to 1+ steelhead trout, the delay in migration of 2+ steelhead trout was most likely due to higher than normal stream discharge during the trapping period in YR 2011.

Similar to 1+ steelhead trout data in YR 2011, average FL and Wt for 2+ steelhead trout in YR 2011 was greater than the average size of previous study years. However, average FL and Wt of 2+ steelhead smolts showed little variation among study years; the greatest difference between any two years was 7.5 mm and 6.0 g. Such differences are unlikely to have biological meaning unless they affect survival to adulthood, which seems doubtful. 2+ steelhead trout are also expected to gain additional length and weight when residing in the estuary prior to ocean entry.

The percentage of 2+ steelhead trout showing parr characteristics was zero each study year, and indicated 2+ steelhead trout do not emigrate through lower Redwood Creek in a parr stage (stream resident form) (author). Rather, most of the 2+ steelhead trout are emigrating in a smolt form. The percentage of 2+ steelhead trout emigrants showing smolt characteristics in YR 2011 (99.3%) was greater than five of the previous study years, however the greatest differences among study years was only six percentage points. In YR 2011, 0.7% of the smolts were classified as pre-smolts, compared to 2.6% for the previous seven year average. Analysis of trapping data ($n = 10$ years) in upper Redwood Creek during YRS 2000 – 2009 showed that smolt percentages in a given study year were negatively related to 2+ steelhead trout population abundances, and negatively related to stream temperatures (Sparkman 2010). Thus, there were less smolt designations for higher population abundances and during study periods with higher stream temperatures. Quinn (2005) reported that stream temperatures play an important role in smoltification, and our data from the upper basin showed that 54% of the variation in smolt percentages over ten study years can be attributed to the variation in stream temperatures (Sparkman 2010). Whether this will be true for 2+ steelhead trout populations emigrating through lower Redwood Creek remains to be tested.

2+ steelhead trout are actively emigrating from upper Redwood Creek through lower Redwood Creek because the trap in lower Redwood Creek (RM 4) has consistently captured efficiency trial fish each study year. Additionally, 2+ steelhead trout from upper Redwood Creek have been observed in the estuary of Redwood Creek every year since

the beginning of our smolt trapping studies (Dave Anderson, pers. comm. 2011). Elastomer marked 2+ steelhead trout released at the upper trap in YRS 2004 and 2005 were also captured by the lower trap in those years. Future trapping efforts will try to increase the sample size of recaptured 2+ steelhead trout for travel time experiments by increasing the sample size of releases from upper Redwood Creek. The lack of large numbers of 3+ steelhead trout provide more evidence that 2+ steelhead trout are actively migrating to the ocean, rather than re-distributing to later migrate to the ocean at age 3.

Although there seems to be few studies that specifically look at steelhead smolt to adult survival, steelhead life history studies in a British Columbia stream (Keogh River) show there is a positive linear relationship between out-migrating 2+ smolts and returning adult steelhead (Ward and Slaney 1988, Ward 2000, Ward et al. 2002). Ward (2000) cites other authors who report similar positive linear relationships between smolts and adults along the British Columbia coast as well (eg Smith and Ward 2000). Survival from smolt to adult in the Keogh River can be variable, and may range from an average of 15% (during 1976-1989) to an average of 3.5% (during 1990-1995) (Ward 2000). Ward and Slaney (1988), reporting on data from the Keogh River for 1978 – 1982 cohorts, determined survival from smolt to adult ranged from 7% to 26%, and averaged 16%. Meehan and Bjornn (1991) reported steelhead smolt to returning adult survival can be a relative high ranging from 10 – 20% in streams that are coastal to a low survival of 2% in streams where steelhead must overcome dams and travel long distances to reach spawning grounds. It is difficult to make specific inferences about 2+ steelhead trout smolt to adult survival for Redwood Creek steelhead based upon successful studies in the literature because of differences in latitude/longitude, geography, ocean conditions (physical and biological), estuaries, and trap locations in the watershed. However, the belief that the number of 2+ smolts relate to future adults (and watershed conditions) is hard to dismiss or invalidate.

With respect to younger juvenile stages (0+ and 1+), the 2+ steelhead trout smolt is the best candidate for assessing steelhead status, trends, and abundance when information on adult steelhead is unavailable or un-attainable. 2+ steelhead trout have overcome the numerous components of stream survival that younger steelhead (0+ and 1+) have not yet completely faced (over-summer, over-winter, etc), and 2+ steelhead smolts are the most direct, juvenile recruit to adult steelhead populations. The 2+ steelhead trout are also an excellent indicator of watershed and stream conditions because they spend the longest amount of time in freshwater habitat prior to ocean entry. Along these same lines, Ward et al. (2003) reported that the 2+ steelhead smolt was a more reliable response variable with respect to stream restoration than late summer juvenile densities because of being less variable.

Cutthroat Trout

A very low number of cutthroat trout were captured in each study year relative to other juvenile salmonids, and over the current eight years a total of 289 individuals were

captured passing through lower Redwood Creek. The total catch of cutthroat trout over eight years was about 99.4% less than the total catch of 1+ and 2+ steelhead trout. The population abundance of cutthroat trout passing through lower Redwood Creek also shows that relatively few cutthroat trout emigrated from the majority of the Redwood Creek basin. The lowest abundance occurred in YR 2008 (N = 54), the highest abundance occurred in YR 2010 (N = 256), and abundance in YR 2011 (N = 118) was the second highest of record. The average abundance over the five years when we determined population abundances equaled 122 individuals. Currently the trend in abundance is non-significantly positive. Low catches and low population abundances does support our hypothesis that few cutthroat trout are emigrating from the majority of the Redwood Creek basin, upstream of the confluence with Prairie Creek. Trap catches in upper Redwood Creek over a twelve year period were even lower, and in YR 2011 only four were captured (Sparkman, In progress). Similar to juvenile coho salmon, the Prairie Creek watershed is probably the biggest contributor to cutthroat trout populations in the Redwood Creek basin based upon this study, the study in upper Redwood Creek, and various studies in Prairie Creek (Walter Duffy, pers. comm. 2011, Duffy and Sparkman, In progress). For example, the smolt trap in lower Prairie Creek (Rm 0.25) in YR 2011 captured 1,198 cutthroat trout smolts, and the resulting population estimate equaled 5,224 individuals (Duffy and Sparkman, In progress).

Similar to other species at age in YR 2011, monthly population emigration through lower Redwood Creek in YR 2011 peaked later (August) than previous study years with exception to YR 2010 when August was the also peak month. The two most important months were July and August in YR 2011 and for the average of the previous years. In Prairie Creek, May was the most important month for population emigration in YR 2011 (Duff and Sparkman, In progress).

All cutthroat trout captured in lower Redwood Creek in YR 2011 were in a smolt stage. An unknown number or percentage of cutthroat trout will residualize in the stream for varying years, and not out-migrate to the estuary and ocean; thus the low trap catches (and population estimates) may not necessarily reflect a very low population size in Redwood Creek. However, if there were large numbers present within the Redwood Creek basin, we would probably catch much more than we do, as they re-distribute or migrate downstream. For example, juvenile salmonid trapping efforts in Prairie Creek consistently capture hundreds of cutthroat trout during spring/early summer as they migrate downstream (Roelofs and Klatt 1996, Roelofs and Sparkman 1999, Walter Duffy, pers. comm. 2011) (Duffy and Sparkman, In progress).

We did not consider any of the young-of-year steelhead trout to be progeny of cutthroat trout because few age-1 and older cutthroat trout were captured in any given year. Far more age 1 and older juvenile steelhead trout (1+ and 2+) migrated through lower Redwood Creek than cutthroat trout as evidenced by trap catches. In the eight study years, for example, the ratio of 1+ and 2+ steelhead trout combined catches to cutthroat trout catches each year ranged from 47:1 to 596:1, and averaged 248:1. In other words there was, on average, 248 times more 1+ and 2+ steelhead trout (combined) captured than cutthroat trout. Ratios would be even higher if population data were used instead of

catch data; thus it seems very unlikely that low numbers of cutthroat trout could produce a significant portion of the young of year trout captures. Therefore, we considered the percentage of 0+ cutthroat trout included in the 0+ steelhead trout catch to be low and negligible.

We used three characteristics to identify coastal cutthroat trout: upper maxillary that extends past the posterior portion of the eye, slash marks on the lower jaws, and hyoid teeth; spotting is also usually more abundant on coastal cutthroat trout. Hybrid juveniles, the product of mating between steelhead trout and cutthroat trout, are commonly noted to be missing one or two of these characters. We have observed less than six potential hybrids in the eight years of study, and based upon visual identification, the number of (potential) hybrids (age-1 and greater) is extremely rare in Redwood Creek. Similar findings occurred in upper Redwood Creek as well (Sparkman 2011b).

0+ Coho Salmon

Similar to 0+ steelhead trout, trap catches of 0+ coho salmon are not all inclusive because only a given percentage of the total number present (upstream of the trapping site) will migrate downstream, this also pertains to the population point estimate. Thus, catches and population estimates are for those fish that were migrating past the trapping site. Trap catches of 0+ coho salmon moving downstream is typical for most streams, including relatively, pristine streams like Prairie Creek (Duffy and Sparkman, In progress). Koski (2009) called these migrating 0+ coho salmon ‘nomads’ and considered this life history strategy to be important for the species resilience.

Few 0+ coho salmon were captured by the trap in lower Redwood Creek in eight consecutive study years (total catch = 1,312 individuals), and in YR 2011 only 226 individuals were captured. The low catches of 0+ coho salmon in lower Redwood Creek is contrasted by high catches in Prairie Creek. For example, trap catches of 0+ coho salmon in mid to upper Prairie Creek from 1996 – 1998 ranged from a low of 372 to a high of 25,492, and averaged 9,659 per trapping season (Roelofs and Sparkman 1999). However, trap catches in lower Prairie Creek in YR 2011 equaled 223 individuals (Duffy and Sparkman, In progress).

In YR 2011 we determined the population abundance of emigrating 0+ coho salmon for the sixth time during our monitoring studies. Population abundances over years equaled 884 in YR 2011, 10 in YR 2010, 63 in YR 2009, 1,886 in YR 2008, 1,057 individuals in YR 2007, and 508 in YR 2006. The preliminary trend in abundance was significantly positive in YR 2008, and with addition of data in YRS 2009 - 2011, showed a non-significant, negative relationship. The current trend line is very susceptible to change due to low sample sizes (n = 6 years), as discussed in earlier sections. Although we may not have a reliable trend in abundance, the point estimates (of abundance) were low each study year, and indicate that relatively few young-of-year coho salmon emigrated through lower Redwood Creek, upstream of the confluence with Prairie Creek. Population

emigration through lower Prairie Creek was also low in YR 2011 (N = 726) (Duffy and Sparkman, In progress).

The migration of 0+ coho salmon at the population level in YR 2011 primarily occurred in the later part of the trapping season, and may be attributable to the increase in stream discharge during the trapping period compared to previous study years. The most important month for migration was July in YR 2011, compared to June for the average of the previous years. Migration in August 2011 was eight times greater than the previous year's average for August. Unlike YR 2010, the temporal delay in migration did not influence the average size of 0+ coho salmon in YR 2011. The average FL and Wt in YR 2011 was slightly greater than the average for the previous seven years, and much less than the average size determined in YR 2010.

The migration of 0+ coho salmon through lower Redwood Creek indicated that these fish were moving downstream to rear, or possibly to enter the ocean at age 0. If the young-of-year coho salmon do not move into Prairie Creek, then they must be moving downstream to the estuary. Thus, lower Redwood Creek and the estuary may serve as an important place for young-of-year coho salmon to rear. Madej et al. (2006) reported that large areas of the Redwood Creek mainstem may be inhospitable for juvenile coho salmon to rear because of high stream temperatures during summer months. Data from trapping efforts in the upper basin support this assertion because juvenile coho salmon were only captured in two of the twelve consecutive study years (Sparkman, In progress). Therefore, determining the spatial distribution of 0+ coho salmon within the Redwood Creek basin is recommended.

1+ Coho Salmon

Low numbers of one plus-year-old coho salmon were caught at the lower trap each study year, with the total catch over eight years equaling 597 individuals. The highest catch occurred in YR 2008 (n = 242), and the lowest catch occurred in YR 2010 (n = 13). Similar to 0+ coho salmon, the low catches of 1+ coho salmon in lower Redwood Creek are contrasted by much higher catches in Prairie Creek. For example, trap catches of 1+ coho salmon in Prairie Creek from 1996 – 1999 ranged from 1,475 – 2,302, and averaged 1,965 per trapping season (Roelofs and Sparkman 1999). More recently, 2,455 1+ coho salmon were captured in lower Prairie Creek in YR 2011 (Duffy and Sparkman, In progress).

The population abundance of 1+ coho salmon in Redwood Creek in YR 2011 was the third lowest of record (N = 113), and about 70% less than the previous seven year average abundance. The average population abundance over eight years of study was 378 individuals, and should be considered to be a very low number. The trend in abundance from YR 2004 - 2011 was non-significantly negative, and indicates more study years will be required to determine a reliable trend in abundance. Population estimates for 1+ coho salmon should be viewed cautiously (due to relatively large error terms for some estimates, 48 - 86%), and the proper context could be that we are 95%

sure that the greatest population abundance during any given study year was less than 1,127 individuals (upper 95% CI for YR 2008 estimate). The population abundances we determined over the current, eight year period can be considered very low (alarmingly so), particularly for a stream the size of Redwood Creek. The low abundances observed in YR 2011 (N = 113), YR 2010 (N = 33), and YR 2007 (N = 102) may indicate year class failures. The population abundance in Prairie Creek in YR 2011 equaled 8,446 individuals, and indicates that Prairie Creek is extremely important for production of coho salmon smolts, and an important refuge for coho salmon in general (Duffy and Sparkman, In progress).

1+ coho salmon in Redwood Creek had the most restricted temporal pattern to migration compared to other juvenile salmonids, with exception to 1+ Chinook salmon. The majority of the 1+ coho salmon population emigrated during May for any given study year, and in YR 2011 May accounted for 90% of the total. The two most important months are usually April and May. Unlike other species at age in YR 2011, 1+ coho salmon did not have delays in migration, and the weekly peak occurred earlier than several study years. The migration of 1+ coho salmon smolts in Prairie Creek in YR 2011 was much more protracted compared to the smolt migration through lower Redwood Creek; the migratory period through lower Prairie Creek in YR 2011 ended on August 6th, compared to an end date of June 21st for lower Redwood Creek.

The average size of 1+ coho salmon smolts in YR 2011 (FL = 107 mm, Wt = 13.40 g) was slightly above average, and close in value to the average size in Prairie Creek in YR 2011 (FL = 109 mm, Wt = 13.47) (Duffy and Sparkman, In progress).

The reason(s) for the lack of sufficient numbers of 1+ coho salmon emigrating from Redwood Creek warrants further study, as does their current distribution within the Redwood Creek basin.

0+ Pink Salmon

Pink salmon in California are recognized as a “Species of Special Concern”, and California is recognized as the most southern border for the species (CDFG 1995). Although not in large numbers, pink salmon have been historically observed in the San Lorenzo River, Sacramento River and tributaries, Klamath River, Garcia River, Ten Mile River, Lagunitas River, Russian River, American River, Mad River, and once in Prairie Creek, which is tributary to Redwood Creek at RM 3.7. Pink salmon were observed spawning in the Garcia River in 1937 and the Russian River in 1955 (CDFG 1995). More recently, adult pink salmon were seen spawning in the Garcia River in 2003 (Scott Monday pers. comm. 2004) and in Lost Man Creek (tributary to Prairie Creek) in 2004 (Baker Holden, pers. comm. 2005). More recently, adult pink salmon were observed and photographed in lower Redwood Creek during the fall of YR 2010 (D. Anderson, pers. com. 2011).

I know of no historic records or anecdotal information documenting pink salmon presence in the mainstem of Redwood Creek prior to our downstream migration trapping efforts. The pink salmon in Redwood Creek are in very low numbers, and were only observed in lower Redwood Creek in YR 2005. In upper Redwood Creek we have captured small numbers of juvenile pink salmon in six out of twelve years, with the most recent capture occurring in YR 2011 (n = 1) (Sparkman, In progress). It is hard to say if the parents of the juvenile pink salmon were strays or remnants of a historic run because so little information exists about adult salmon in Redwood Creek. According to the Habitat Conservation Planning Branch (HCPB) of CDFG, pink salmon are considered to be “probably extinct” in California (CDFG 1995). However, the HCPB does state that “more efforts need to be conducted to prove (or disprove) that reproducing populations exist anywhere in California” (CDFG 1995). Based upon our trapping data in upper and lower Redwood Creek, pink salmon are present in Redwood Creek and reproducing, albeit in low numbers.

CONCLUSIONS

The migration of juvenile salmonids through lower Redwood Creek consisted of juvenile Chinook salmon (ocean-type and to a much lesser degree stream-type), steelhead trout (at least three age classes), coho salmon (two age classes), and cutthroat trout (multiple age classes, including adults). The abundance of 0+ Chinook salmon in YR 2011 (N = 147,719) was greater than YRS 2005 – 2007 and YR 2010, and 28% less than the previous seven year average. The abundance of 1+ Chinook salmon in YR 2011 equaled 15 individuals, and indicated for the third year in a row that yearling Chinook salmon are relatively rare in Redwood Creek. The abundance of 1+ steelhead trout in YR 2011 (N = 20,501) was the lowest of record, and for 2+ steelhead trout (N = 5,587) was the second lowest of record. Far more 1+ steelhead trout emigrated from Redwood Creek than 2+ steelhead trout each year, and may indicate stream habitat conditions are limiting the abundance of the older age class (2 years); or favoring a change in life history to a younger smolt age (age 1). The abundance of 0+ coho salmon in YR 2011 was a low value of 884 individuals, however, not all of the 0+ coho salmon within the Redwood Creek basin will migrate downstream. The 0+ coho salmon migrating through lower Redwood Creek (and Prairie Creek) are no longer considered ‘lost or surplus production’, and several studies are showing this life history can give reliance to populations. The abundance of 1+ coho salmon smolts passing through lower Redwood Creek was a very low value of 113 individuals, and may indicate a cohort failure (for the population upstream of the confluence with Prairie Creek) for the second year in a row. The abundance of cutthroat trout in YR 2011 (N = 118) was the second highest of record, however, abundances across all years can be considered to be low.

The trends in population abundances over the current eight years were non-significant for 0+ Chinook salmon, 0+ coho salmon, 1+ coho salmon, and cutthroat trout. The trends in abundance for 1+ steelhead trout smolts and 2+ steelhead trout smolts over time were significantly negative. The addition of whether or not there were flood type flows in the upper basin during winter months was an important, additional variable for analyzing the

trends of 0+ Chinook salmon and 1+ steelhead trout. The addition of flood type flows in the model for 0+ Chinook salmon caused the trend over time to become statistically significant. With respect to 0+ Chinook salmon, population abundances were negatively related to study year and flood type flows, and indicated that the 0+ Chinook salmon populations passing through lower Redwood Creek were strongly tied to abundances in upper Redwood Creek and flood type flows in the upper basin which occurred during the redd egg incubation/embryogenesis periods. For both 1+ steelhead trout and 2+ steelhead trout, abundances were negatively related to study year and flood type flows in the upper basin, which may indicate that over winter conditions in the upper basin are limiting survival when flood type flows are present.

The patterns in monthly population abundances showed a temporal delay in migration for several species at age which I attribute to increased stream discharge during the trapping period, compared to previous study years. Trapping in YR 2011 was extended into September to account for delays in migration, and delays in migration were readily apparent for 0+ Chinook salmon, 1+ Chinook salmon, 1+ steelhead trout, 2+ steelhead trout, 0+ coho salmon, and cutthroat trout. The two most important months for migration were June and July for 0+ Chinook salmon, 1+ steelhead trout, 2+ steelhead trout, and 0+ coho salmon; May and June for 1+ Chinook salmon, April and May for 1+ coho salmon, and July and August for cutthroat trout.

The trap catch of 0+ steelhead trout in YR 2011 ($n = 9,864$) was the fourth lowest of record, which could be attributable to: 1) an decrease in adult numbers upstream of the trap site, 2) lower trapping efficiencies, 3) difference in the percentage of the total 0+ steelhead trout population emigrating downstream each year, or 4) some combination of factors 1 – 3. Marked 0+ steelhead trout released at the upper trap in Redwood Valley in YRS 2006, 2007, 2010, and 2011 were recaptured at the lower trap in the same year as the release, and indicate 0+ steelhead trout can migrate considerable distances in search of rearing areas. These experiment could be the first to document long range dispersal (29 mi.) of young of year steelhead trout from spawning to rearing areas. The 0+ steelhead trout and 0+ coho salmon that passed by the trap in lower Redwood Creek are probably rearing in reaches below RM 4 and in the estuary, thus lower Redwood Creek and the estuary are also important for young-of-year fish, in addition to older, juvenile salmonid age classes.

The overall study objectives in YR 2011 were successfully completed, and demonstrated that the smolts passing through lower Redwood Creek can be monitored, even under higher than average stream flows. During some years we may fail in obtaining population abundance estimates for cutthroat trout or 0+ coho salmon due to low sample sizes, however, we may be able to model the data using data from previous study years (i.e. modeling trapping efficiency with stream flow) or by using trapping efficiencies associated with fish of nearly equal size and age in the same study year.

Redwood Creek is currently classified as temperature and sediment impaired, and many scientists, resource managers, and members of the public also realize that the estuary is in need of restoration. Future fisheries work in Redwood Creek should be able to show any

fish response to current and future watershed and stream conditions by combining data from this study, smolt trapping in upper Redwood Creek, adult and juvenile studies in Prairie Creek, and juvenile monitoring in the estuary (author, Walter Duffy pers. comm. 2011, and David Anderson, pers. comm. 2011). In addition, the Department of Fish and Game and USGS CA. Cooperative Fish and Wildlife Research Unit have begun monitoring adult salmon and steelhead populations in Redwood Creek; if possible, such information could be used to investigate smolt to adult, and adult to smolt relationships. The adult information would greatly compliment our studies on smolt populations within the Redwood Creek basin.

RECOMMENDATIONS

This study is one of the few studies that is designed to document smolt abundance and population trends of the California Coastal Chinook salmon ESU, Southern Oregon/Northern California Coasts Coho salmon ESU, Northern California Steelhead Trout ESU, and Southern Oregon/California Coasts Coastal Cutthroat Trout ESU over a relatively long time period. With respect to the Chinook salmon ESU and steelhead trout ESU, this study might be the only one that provides population data for a relatively large stream. The most important recommendation to make is to continue this study over multiple consecutive years (10+) in order to:

1. Encompass as much environmental and biological variation as possible.
2. Cover multiple cohort life cycles over time.
3. Collect baseline data for future comparisons:
 - a. Collect data on juvenile salmonid life histories in Redwood Creek, which will increase our understanding of juvenile salmonids (smolts).
 - b. Detect changes in population abundance which can be used to assess the status and trends of Chinook salmon, steelhead trout, coho salmon, and (possibly) cutthroat trout in Redwood Creek.
 - c. Detect any fish response (population, fish size, age class composition, etc) to stream and watershed conditions and restoration activities in Redwood Creek.
4. Help focus habitat restoration efforts and needs in the basin.
5. Investigate relationships between the number of adults (via redd counts and adult counts using DIDSON technology) and smolt production.

This study, when combined with juvenile salmonid monitoring in the upper basin (RM 33, and estuary (Redwood National Park), will also help determine bottlenecks to anadromous salmonid production in Redwood Creek.

ACKNOWLEDGEMENTS

I thank the California Department of Fish and Game AFRAMP, and the Fisheries Restoration Grants Program (Project Numbers P0810509 and S02067) for funding this study. I would also like to thank the USGS California Cooperative Fish and Wildlife Research Unit (Dr. Walter Duffy, Leslie Farrar), Humboldt State University Sponsored Programs Foundation, and CDFG (Scott Monday) for their assistance in managing the grant contract. I am particularly thankful to Joe Hufford for access to his property, and for his continued interest in salmon and steelhead trout in Redwood Creek. I thank Don Chapman for his interest in the study and for his helpful comments, and Catherine Stone for her general support. Lastly, and by no means in the least, I thank the dedicated field crew (Becky Dutra, Rod Park, Melissa Gilbert, Dominic Vitali, Chase Flowers, Tom Herman, Paula Whitten, Scott Lord, Laurel Osborn, Jasper Amir, and Matt Settelmayer) who counted and examined every fish we captured and marked, for removing debris from the livebox in the evenings and nights, for releasing marked fish upstream of the trap site at night, for helping me monitor the trap during high flow events, and for their positive attitudes.

LITERATURE CITED

- Bendock T. 1995. Marking juvenile Chinook salmon in the Kenai River and Deep Creek, Alaska, 1993-1994. Alaska Department of Fish and Game, Fishery Data Series No. 95-17, Anchorage. 34 p.
- Bradford MJ, RA Myers, and JR Irvine. 2000. Reference points for coho salmon (*Oncorhynchus kisutch*) harvest rates and escapement goals based on freshwater production. Canadian Journal of Fisheries and Aquatic Sciences 57:677-686.
- Brown R. 1988. Physical rearing habitat for anadromous salmonids in the Redwood Creek basin, Humboldt County, California. MS thesis, Humboldt State University, Arcata, California. 132 p.
- Busby PJ, TC Wainwright, GJ Bryant, LJ Lierheimer, RS Waples, FW Waknitz, and IV Lagomarsino. 1996. Status review of west coast steelhead from Washington, Idaho, Oregon, and California. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-NWFSC-27, 261 p.
- Cannata S, R Henly, J Erler, J Falls, D McGuire, and J Sunahara. 2006. Redwood Creek watershed assessment report. Coastal Watershed Planning and Assessment Program and North Coast Watershed Assessment Program. California Resources

- Agency and California Environmental Protection Agency, Sacramento, CA. 166 p.
- Carlson SR, LG Coggins Jr., and CO Swanton. 1998. A simple stratified design for mark-recapture estimation of salmon smolt abundance. *Alaska Fishery Research Bulletin* 5(2):88-102.
- Cashman SM, HM Kelsey, and DR Harden. 1995. Geology of the Redwood Creek Basin, Humboldt County, California. Pages B1 – B13 *in* KM Nolan, HM Kelsey, and DC Marron, editors. *Geomorphic Processes and Aquatic Habitat in the Redwood Creek Basin, Northwestern California*. US Geologic Survey Professional Paper 1454.
- [CDFG] California Department of Fish and Game. 1995. Habitat Conservation Planning Branch, Fish Species of Special Concern in California, Pink Salmon. http://www.dfg.ca.gov/hcpb/cgi-bin/read_one.asp?specy=fish&idNum=58
- [CDFG NCWAP] California Department of Fish and Game. 2004. North Coast Watershed Assessment Program, <http://www.ncwatershed.ca.gov/>
- Cederholm, C.J., L.M. Reid, and E.O. Salo. 1981. Cumulative effects of logging road sediment on salmonid populations in the Clearwater River, Jefferson County, Washington. Contribution No. 543, College of Fisheries, University of Washington, Seattle, Washington. 35 p.
- [CWA] California Water Act. 2002. CWA section 303(d) list of water quality limited segment; approved by the State Water Resources Control Board, February 4, 2003.
- Cleary JS. 2001. Increasing coho productivity in the Chilliwack River Watershed. Pages 1 – 6 *in* Watershed Restoration Technical Bulletin Streamline, Vol. 6 No.1, 20 p.
- Conner WP, JG Sneva, KF Tiffan, RK Steinhorst, and D Ross. 2005. Two alternative juvenile life history types for fall Chinook salmon in the Snake River Basin. *Transactions of the American Fisheries Society* 134:291-304.
- Devries P. 1977. Riverine salmonid egg burial depths: review of published data and implications for scour studies. *Canadian Journal of Fisheries and Aquatic Sciences* 54:1685-1698.
- Ebersole, JL, ME Colvin, PJ Wigington, SG Leibowitz, JP Baker, MR Church, JE Compton, BA Miller, MA Cairns, BP Hansen, HR LaVigne. 2009. Modeling stream network-scale variation in coho salmon overwinter survival and smolt size. *Transactions of the American Fisheries Society* 138:564-580.

- Federal Register. 1997. Endangered and Threatened Species: Threatened status for coho salmon in the Southern Oregon/Northern California Coast Evolutionarily Significant Unit in California. Federal Register, Washington D.C. 62: 24588 – 24609.
- Federal Register. 1999a. Endangered and Threatened Species: Threatened status for two Chinook Salmon Evolutionarily Significant Unit in California. Final Rule. Federal Register, Washington D.C. 64: 50933 – 50415.
- _____ 1999b. Endangered and Threatened Status for Southwestern Washington/Columbia River Coastal cutthroat trout in Washington and Oregon, and delisting of Umpqua River cutthroat trout in Oregon: Proposed Rule. Federal Register, Washington D.C., 64: pps 16397-16413.
- Federal Register. 2000. Endangered and Threatened Species: Threatened status for one steelhead Evolutionarily Significant Unit in California. Federal Register, Washington D.C. 65: 36704 – 36094.
- Fitzgerald JL, TF Sheehan, and JF Kocik. 2004. Visibility of visual implant elastomer tags in Atlantic Salmon reared for two years in marine net-pens. North American Journal of Fisheries Management 24: 222-227.
- Gangmark HA, and RG Bakkala. 1960. A comparative study of unstable and stable (artificial channel) spawning streams for incubating king salmon at Mill Creek. California Fish and Game. 46:151-164.
- Giannico GR, and SG Hinch. 2003. The effect of wood and temperature on juvenile coho salmon winter movement, growth, density, and survival in side-channels. River Research and Applications 19:219-231.
- Greene CM, DW Jensen, GR Pess, EA Steel, and E Beamer. 2005. Effects of environmental conditions during stream, estuary, and ocean residency on Chinook salmon return rates in the Skagit River, Washington. Transactions of the American Fisheries Society 134:1562-1581.
- Hartman GF, and JC Scrivener. 1990. Impacts of forest practices on a coastal stream ecosystem, Carnation Creek, British Columbia. Canadian Bulletin of Fisheries and Aquatic Sciences. 223 p.
- Hicks BJ, JD Hall, PA Bisson, and JR Sedell. 1991. Responses of salmonids to habitat changes. Pages 483 – 518 *in* WR Meehan, editor. Influences of forest and rangeland management on salmonid fishes and their habitats. American Fisheries Society Special Publication 19.
- Hintze J. 1998. Number Crunchers Statistical System. NCSS 97, Version 6.0.

- Holtby LB, and MC Healey. 1986. Selection for adult size in female coho salmon (*Oncorhynchus kisutch*). Canadian Journal of Fisheries and Aquatic Sciences 43:1948-1959.
- Johnson S. 2004. Summary of habitat and fish monitoring data from East Fork and Upper Mainstem Lobster Creeks 1988: 2004. Western Oregon Research and Monitoring Program, Oregon Department of Fish and Wildlife, 13 p.
- Kelsey HM, M Coghlan, J Pitlick, and D Best. 1995. Geomorphic analysis of streamside landslides in the Redwood Creek Basin, Northwestern California. Pages J1 - J12 in KM Nolan, HM Kelsey, and DC Marron, editors. Geomorphic Processes and Aquatic Habitat in the Redwood Creek Basin, Northwestern California. US Geologic Survey Professional Paper 1454.
- Koski, KV. 2009. The fate of coho salmon nomads: the story of an estuarine-rearing strategy promoting resilience. Ecology and Society, 14(1):4, 15 p.
- Madej MA, and V Ozaki. 1996. Channel response to sediment wave propagation and movement, Redwood Creek, California, USA. Earth Surface Processes and Landforms, Vol. 21, 911 – 927.
- Madej MA, C Currens, V Ozaki, J Yee, and DG Anderson. 2006. Assessing possible thermal rearing restrictions for juvenile coho salmon (*Oncorhynchus kisutch*) through thermal infrared imaging and in-stream monitoring, Redwood Creek, California. Canadian Journal of Fisheries and Aquatic Sciences 63:1394-1396.
- Maher FP and PA Larkin. 1955. Life history of the steelhead trout of the Chilliwack River, British Columbia. Transactions of the American Fisheries Society 84:27-38.
- McCubbing D. 2002. Adult steelhead trout and salmonid smolt migration at the Keogh River, B.C. during Spring 2002. Habitat Conservation Trust Fund. Contract Number: CBIO3006. Province of British Columbia, Ministry of Water, Land, and Air Protection, Biodiversity Branch, Fisheries Research and Development, University of British Columbia, Vancouver, BC. 34 p.
- McCubbing DJF, and BR Ward. 1997. The Keogh and Waukwaas rivers paired watershed study for B.C.'s Watershed Restoration Program: juvenile salmonid enumeration and growth 1997. Province of British Columbia, Ministry of Environment, Lands and Parks, and Ministry of Forests. Watershed Restoration Project Report No. 6:33p.
- McCubbing D, and B Ward. 2003. Adult steelhead trout and salmonid smolt migration at the Keogh River, B.C., during winter and spring 2003. Habitat Conservation Trust Fund, Contract Number: CBIO4051. Province of British Columbia, Ministry of

- Water, Land, and Air Protection, Biodiversity Branch, Aquatic Ecosystem Science Section, Vancouver, B.C. 51 p.
- McNeil WH. 1966. Effect of the spawning bed environment on reproduction of pink and chum salmon. Fishery Bulletin Volume 65, Number 2. Contribution Number 198. College of Fisheries. University of Washington, Seattle.
- Meehan WR, and TC Bjornn. 1991. Salmonid distributions and life histories. Pages 47-82 in W.R. Meehan (ed.). Influences of forest and rangeland management on salmonid fishes and their habitats. Special Publication 19, American Fisheries Society. Bethesda, MD.
- Montgomery DR, JM Buffington, NP Peterson, D Schuett-Hames, and TP Quinn. 1996. Streambed scour, egg burial depths, and the influence of salmonid spawning on bed surface mobility and embryo survival. Canadian Journal of Fisheries and Aquatic Sciences 53:1061-1070.
- Murphy ML and WR Meehan. 1991. Stream ecosystems. Pages 17-46. in WR Meehan (ed). Influences of forest and rangeland management on salmonid fishes and their habitats. Special Publication 19, American Fisheries Society. Bethesda, MD.
- NCSS. Number Crunchers Statistical Software. 1997. Jerry Hintze.
- Nickelson TE. 1986. Influences of upwelling, ocean temperature, and smolt abundance on marine survival of coho salmon (*Oncorhynchus kisutch*) in the Oregon production area. Canadian Journal of Fisheries and Aquatic Sciences 43:527-535.
- Newby NC, TR Binder, and ED Stevens. 2007. Passive integrated transponder tags (PIT) tagging did not negatively affect the short-term feeding behavior or swimming performance of juvenile rainbow trout. Transactions of the American Fisheries Society 136:341-345.
- [NOAA] National Oceanic and Atmospheric Administration. 1999. Fact Sheet: West Coast Coho Salmon (*Oncorhynchus kisutch*). 2 p.
- Pautzke CF, and RC Meigs. 1941. Studies on the life history of the Puget Sound steelhead trout (*Salmo Gairdnerii*). Transactions of the American Fisheries Society, 70:209-220.
- Quinn TP, and NP Peterson. 1996. The influence of habitat complexity and fish size on over-winter survival and growth of individually marked juvenile coho salmon (*Oncorhynchus kisutch*) in Big Beef Creek, Washington. Canadian Journal of Fisheries and Aquatic Sciences 53:1555-1564.

- Quinn TP. 2005. The behavior and ecology of pacific salmon and trout. American Fisheries Society, Bethesda Maryland in association with University of Washington Press, Seattle, 378 p.
- [RNP] Redwood National and State Parks. 1997. Redwood Creek Watershed Analysis. In house, 91 p.
-
- _____ . 2007. Redwood Creek Watershed Analysis. In house data.
- Roelofs TD, and B Klatte. 1996. Anadromous salmonid escapement and downstream migration studies in Prairie Creek, California, 1995-1996. Report for the California Department of Transportation, Sacramento, California. 18 p.
- Roelofs TD, and MD Sparkman. 1999. Effects of sediments from the Redwood National Park bypass project (CALTRANS) on anadromous salmonids in Prairie Creek State Park 1995-1998. Final report to the California Department of Transportation, contract No. 001A0162. Department of Fisheries, Humboldt State University, Arcata, California. 28 p.
- Roni P, SA Morley, P Garcia, C Detrick, D King and E Beamer. 2006. Coho salmon smolt production from constructed and natural floodplain habitat. Transactions of the American Fisheries Society 135:1398-1408.
- Roper BB, and DL Scarnecchia. 1999. Emigration of age-0 Chinook salmon (*Oncorhynchus tshawytscha*) smolts from the upper South Umpqua River basin, Oregon, U.S.A. Canadian Journal of Fisheries and Aquatic Sciences 56: 939-946.
- Rowe TA. 2003. Downstream Migrant Trapping Report: Horse Linto Creek and Willow Creek. U.S.D.A. Forest Service, Lower Trinity Ranger Station, Willow Creek, California, USA.
- Schuett-Hames DE, NP Peterson, R Conrad, and TP Quinn. 2000. Patterns of gravel scour and fill after spawning by chum salmon in a Western Washington stream. North American Journal of Fisheries Management 20:610-617.
- Seiler DS, S Neuhauser, and L Kishimoto. 2003. 2002 Skagit River wild 0+ Chinook production evaluation annual report. Washington Department of Fish and Wildlife, Olympia.
- Seiler D, G Volkhardt, P Topping, and L Kishimoto. 2004. 2001 Green River juvenile salmonid production evaluation. Washington Department of Fish and Wildlife, Fish Program, Science Division, Olympia, Washington 98501-1091; 36 p.
- Sharma R, and R Hilborn. 2001. Empirical relationships between watershed characteristics and coho salmon (*Oncorhynchus kisutch*) smolt abundance in 14

- western Washington streams. *Canadian Journal of Fisheries and Aquatic Sciences* 58(7):1453-1463.
- Slaney PA, CJ Perrin, and BR Ward. 1986. Nutrient concentration as a limitation to steelhead smolt production in the Keogh River. *Proceedings of the Annual Conference Western Association of Fish and Wildlife Agencies* 66:146-158.
- Smith BD, and BR Ward. 2000. Trends in wild adult steelhead (*Oncorhynchus mykiss*) abundance for coastal regions of British Columbia support the variable marine survival hypothesis. *Canadian Journal of Fisheries and Aquatic Sciences* 57:271-284.
- Solazzi MF, TE Nickelson, SL Johnson, and JD Rodgers. 2000. Effects of increasing winter rearing habitat on abundance of salmonids in two coastal Oregon streams. *Canadian Journal of Fisheries and Aquatic Sciences* 57:906-914.
- Sparkman, MD. 2002. Mad River juvenile steelhead downstream migration study, Humboldt County, Ca. CDFG S-RAMP, 2002 Annual report, project 2a3:32 p.
- _____ 2010. Upper Redwood Creek juvenile salmonid (smolt) downstream migration study, study year 2008. California Department of Fish and Game, Anadromous Fisheries Resource Assessment and Monitoring Program, 2008 Annual Report, 2a5: 131 p.
- _____ 2011a. Lower Redwood Creek juvenile salmonid (smolt) abundance project, study year 2009. California Department of Fish and Game, Anadromous Fisheries Resource Assessment and Monitoring Program, 2009 Annual Report, 2a7: 138 p.
- _____ 2011b. Upper Redwood Creek juvenile salmonid (smolt) abundance project, study year 2010: a report to the Fisheries Restoration Grants Program. California Department of Fish and Game, Anadromous Fisheries Resource Assessment and Monitoring Program, study 2a5: 51 p.
- [SWRCB]. State Water Resources Control Board. 2003. Resolution number 2003 – 0009. Approval of the 2002 Federal Clean Water Act Section 303 (d) List of Water Quality Limited Sections.
- Teel DJ, GB Milner, GA Winans, and WS Grant. 2000. Genetic population structure of life history types in Chinook Salmon in British Columbia, Canada. *Transactions of the American Fisheries Society* 129:194-209.
- Tripp DB, and VA Poulin. 1986. The effects of logging and mass wasting on salmonid spawning habitat in streams on the Queen Charlotte Islands. British Columbia Ministry of Forests and Lands, Land Management Report 50, Vancouver.

- Tripp DB. 1986. Using large organic debris to restore fish habitat in debris-torrented streams. British Columbia Ministry of Forests and Lands, Land Management Report 47, Victoria.
- Unwin MJ. 1997. Fry-to-adult survival of natural and hatchery-produced Chinook salmon (*Oncorhynchus tshawytscha*) from a common origin. Canadian Journal of Fisheries and Aquatic Sciences 54:1246-1254.
- [USEPA] United States Environmental Protection Agency. 2003. Approval of the Clean Water Act Section 303(d) List of Water Quality Limited Segments as approved by the State Water Resources Control Board on February 4, 2003 as Resolution No. 2003-0009.
- [USFWS] United States Fish and Wildlife Service. 2001. Juvenile salmonid monitoring on the mainstem Klamath River at Big Bar and mainstem Trinity River at Willow Creek, 1997 – 2000. Annual Report of the Klamath Fisheries Assessment Program, Arcata Fish and Wildlife Office, Arcata, CA. 106 p.
- [USGS] United States Geological Survey. 2010. Surface water data for California: monthly streamflow statistics. Web page http://waterdata.usgs.gov/ca/nwisdv?cb_00060=on&format=rdb&begin_date=2008-03-15&end_date=2009-03-15&site_no=11482500&referred_module=sw
- Ward BR. 2000. Declivity in steelhead (*Oncorhynchus mykiss*) recruitment at the Keogh River over the past decade. Canadian Journal of Fisheries and Aquatic Sciences 57:298-306.
- Ward BR, DJF McCubbing, and PA Slaney. 2003. The addition of inorganic nutrients and stream habitat structures in the Keogh River Watershed for steelhead trout and coho salmon. Pages 127–147 in JG Stockner, (ed.). Proceedings of the International Conference on Restoring Nutrients to salmonid ecosystems, April 24 – 26, 2001, Eugene. Oregon.
- Ward BR, DJF McCubbing, and PA Slaney. 2002. Stream restoration for anadromous salmonids by the addition of habitat and nutrients. Sixth International Atlantic salmon Symposium, 15th – 18th July, 2002. Edinburgh, Scotland. 23 p.
- Ward BR, and PA Slaney. 1988. Life history and smolt-to-adult survival of Keogh River steelhead trout (*Salmo gairdneri*) and the relationship of smolt size. Canadian Journal of Fisheries and Aquatic Sciences 45:1110-1122.
- Ward BR, PA Slaney, AR Facchin, and RW Land. 1989. Size-based survival in steelhead trout (*Oncorhynchus mykiss*): back-calculated lengths from adults' scales compared to migrating smolts at the Keogh River, British Columbia. Canadian Journal of Fisheries and Aquatic Sciences Vol. 46, pgs 1853-1858.

Zar JH. 1999. Biostatistical Analysis. Prentice Hall, Upper Saddle River, New Jersey. 663 p.

PERSONAL COMMUNICATIONS

Anderson, David. 2000 - 2011. Redwood National and State Parks, Fisheries Biologist, Orick, California.

Chapman, Don W. 2000 - 2011. McCall, Idaho.

Chesney, Bill. 2006. California Department of Fish and Game, Associate Fisheries Biologist, Yreka, Ca.

Duffy, Walter G. 2005 - 2011. California Cooperative Fishery Research Unit Project Leader, U.S. Geological Survey, Humboldt State University, Arcata, CA.

Holden, Baker. 2005. Redwood National and State Parks, Fisheries Biologist Orick, CA.
Madej, Mary Ann. 2005 - 2011. United States Geological Survey, Research Geologist, Arcata, CA.

Monday, Scott. 2004. California Department of Fish and Game, Fortuna, CA.

Ozaki, Vicki. 2004 - 2011. Redwood National Park Service, Geologist. Arcata, CA. 95519.

Peters, Charlotte. 2001. Redwood Creek Watershed Map. Information Services, California Department of Fish and Game, Eureka, CA.

Stover, Marlin. 2000. Upper Redwood Creek long time resident and property owner. Redwood Valley, Humboldt County, California.

APPENDICES

Appendix 1. Reasons for collecting genetic samples from Chinook salmon, steelhead trout smolts, and coho salmon fry, parr, and smolts.

Chinook Salmon:

1. To test for possible genetic differences between 0+ Chinook (Ocean-Type) and 1+ Chinook (Stream-Type).
2. To test for possible genetic differences between 0+ Chinook salmon fry and 0+ Chinook salmon fingerlings.

Steelhead Trout:

1. To test for any hatchery introgression into the wild steelhead stock in Redwood Cr.
2. To test for possible genetic differences between age-1 and age-2 smolts.
3. To test for possible genetic differences between emigrating 0+ steelhead trout and 1+ steelhead trout the following year.

Coho Salmon

1. To determine the number of parents responsible for the juveniles captured in the fish trap.

All Species:

1. To test for possible genetic differences between fish captured in the lower basin and upper basin.
2. To construct a genetic data base for future comparisons and analyses.

Appendix 2. Graphical representation of daily stream gage height (ft.) at trap site and (preliminary) average daily streamflow (cfs) measured at Orick gaging station (USGS 2011), lower Redwood Creek, Humboldt County, CA.

